

Control of electron spin decoherence caused by electron–nuclear spin dynamics in a quantum dot

Ren-Bao Liu^{1,2}, Wang Yao¹ and L J Sham^{1,3}

¹ Department of Physics, University of California, San Diego, La Jolla, CA 92093-0319, USA

² Department of Physics, The Chinese University of Hong Kong, Shatin, NT, Hong Kong, People's Republic of China

E-mail: rbliu@phy.cuhk.edu.hk and lsham@ucsd.edu

New Journal of Physics **9** (2007) 226

Received 28 March 2007

Published 11 July 2007

Online at <http://www.njp.org/>

doi:10.1088/1367-2630/9/7/226

Abstract. Control of electron spin decoherence in contact with a mesoscopic bath of many interacting nuclear spins in an InAs quantum dot is studied by solving the coupled quantum dynamics. The nuclear spin bath, because of its bifurcated evolution predicated on the electron spin up or down state, measures the which-state information of the electron spin and hence diminishes its coherence. The many-body dynamics of the nuclear spin bath is solved with a pair-correlation approximation. In the relevant timescale, nuclear pair-wise flip–flops, as elementary excitations in the mesoscopic bath, can be mapped into the precession of non-interacting pseudo-spins. Such mapping provides a geometrical picture for understanding the decoherence and for devising control schemes. A close examination of nuclear bath dynamics reveals a wealth of phenomena and new possibilities of controlling the electron spin decoherence. For example, when the electron spin is flipped by a π -pulse at τ , its coherence will partially recover at $\sqrt{2}\tau$ as a consequence of quantum disentanglement from the mesoscopic bath. In contrast to the re-focusing of inhomogeneously broadened phases by conventional spin-echoes, the disentanglement is realized through shepherding quantum evolution of the bath state via control of the quantum object. A concatenated construction of pulse sequences can eliminate the decoherence with arbitrary accuracy, with the nuclear–nuclear spin interaction strength acting as the controlling small parameter.

³ Author to whom any correspondence should be addressed.

Contents

1. Introduction	2
2. Theory	6
2.1. The model	6
2.2. General formalism	9
2.3. Essential assumptions	10
2.4. Ensemble and single-system dynamics	11
2.5. Pair-correlation approximation and pseudo-spin model for nuclear spin dynamics	14
3. Free-induction decay	16
4. Hahn echo signal	22
5. Disentanglement and recoherence	27
6. Concatenated disentanglement	31
6.1. Formalism and geometrical picture	31
6.2. Short-time behaviours	33
6.3. Controlling small parameters	35
6.4. Coherence stabilization	37
7. Summary and discussions	37
Acknowledgments	40
Appendix A. Factorization of single-system dynamics and inhomogeneous broadening	40
Appendix B. Boundary of mesoscopic nuclear spin bath	40
Appendix C. Pair-flip numbers	41
Appendix D. Error estimation for pair-correlation approximation	43
References	45

1. Introduction

A quantum system, unlike a classical one, can be in a coherent superposition of constituent states. This coherence is the wellspring of quantum properties and is key to quantum technology. The contact of a quantum object with a macroscopic system causes loss of state coherence [1]–[3]. Advances towards quantum technology have effectively substituted the macroscopic environments by mesoscopic ones. Usually, the contact interaction between the quantum object and a ‘particle’ of the bath weakens with increasing bath size (defined by the number of particles, N), while the interaction between particles inside the bath is independent of the bath size. When the bath size is in the mesoscopic regime such that the object–bath interaction dominates their interaction with the rest of universe, the quantum object and the mesoscopic bath evolve as a closed system in the relevant timescales. Thus, a proper description of the quantum object in such a mesoscopic bath is given by the full quantum mechanical solution of the coupled object–bath evolution [4]–[9], as opposed to the semiclassical treatment in conventional decoherence studies [10, 11].

Spins of single electrons confined in semiconductor quantum dots (QDs) are paradigmatic systems in mesoscopic physics [12, 13] and in spin-based quantum technology [14]–[16]. Spin

decoherence is a main limiting factor to quantum properties and has been extensively studied both in theories [4]–[9], [17]–[30], and in experiments [31]–[41]. It has been well established by theories [27]–[30] and experiments [31]–[34] that the electron spin decoherence caused by phonon scattering is negligible at a temperature lower than a few Kelvin. The relevant bath at low temperature then is a spin bath [42] composed of lattice nuclear spins in a QD. The electron spin is coupled to a nuclear spin through the contact hyperfine interaction with magnitude inversely proportional to the total number N of nuclei in the QD. For the QD size of interest, the hyperfine coupling is much stronger than the mutual interaction between nuclear spins. Therefore, a mesoscopic bath consisting of all nuclear spins within the QD (i.e. in direct contact with the electron) is identified. Electron spin decoherence at low temperatures is determined by the quantum dynamics of the mesoscopic spin system.

Conventionally, the relaxation of spin coherence (or spin polarization) is categorized into two types, namely, the T_1 (longitudinal) relaxation and the T_2 (transverse) relaxation. Regarding the density matrix in the basis of the energy eigenstates, the T_1 process causes the redistribution of diagonal elements accompanied by the decay of off-diagonal ones, and the T_2 process causes the decay of off-diagonal elements. In ensemble measurements, the static random distribution of external fields (i.e. inhomogeneous broadening) also leads to an effective dephasing or T_2^* process. In [2], the T_2^* process is called ‘fake’ decoherence as it results from the classical phase average, the dephasing in the T_1 process is called ‘false’ decoherence as it is due to the spin population redistribution, and only the T_2 process excluding the effect accompanying the spin flip (often called pure dephasing) is dubbed the ‘true’ decoherence since essentially it originates from quantum entanglement between the system and the bath. In the specific electron–nuclear spin system considered here, the T_2^* process is due to the thermal distribution of the Overhauser field from the nuclear spins [17, 21]. The T_1 process is due to the off-diagonal hyperfine interaction [17, 21, 24, 27]. The T_2 process is because of the electron–nuclear spin entanglement established during the nuclear spin evolution driven by the hyperfine interaction and the nuclear–nuclear spin interaction [43]. As the hyperfine interaction is the strongest part of the spin interaction in a QD, initial theoretical efforts have been devoted to the spin decoherence by the hyperfine interaction only where the (intrinsic) nuclear–nuclear spin interaction is neglected [17]–[24]. Such theories are successful in understanding many essential parts of the spin decoherence in QDs, especially the T_2^* and T_1 processes. But there are important situations under which we have to take into account the intrinsic nuclear–nuclear spin interaction (the term ‘intrinsic’ is used to contrast the ‘extrinsic’ interaction between nuclei mediated by the hyperfine interaction). Firstly, under a moderate to strong magnetic field ($B \gtrsim 0.1$ T), the Zeeman energy mismatch between the electron and the nuclei will suppress the spin relaxation effect of the hyperfine interaction (including both T_1 and T_2 processes) [4, 21, 22, 24] and hence the nuclear–nuclear spin interaction emerges to be important. Secondly and more importantly, it has been shown that the leading order effect of the hyperfine interaction in the dephasing is removed by a spin echo [4, 23] and thus spin echo signals would not be correctly accounted for without considering the intrinsic interaction. The pure dephasing effect of the intrinsic nuclear spin interaction in QD was first studied in a semiclassical framework [25, 26] which gives correct timescales of the decoherence but misses several important features such as the exponential index of the decay profile (especially in the pulse control configurations [44]). Recently, quantum theories including the intrinsic interaction have been developed independently by the Das Sarma group [7]–[9] and the Sham group [4]–[6]. The two theories agree with each other in the lowest level of approximation (i.e. the pair-correlation approximation as discussed below). The Sham group’s linked cluster expansion

method [6] provides a systematic method to include the higher-order correlations order by order, which does not agree completely with the density matrix cluster expansion theory by the Das Sarma group [9]. We focus on the T_2 decoherence caused by the electron–nuclear entanglement in which the nuclear–nuclear spin interaction is essential. To separate the quantum entanglement effect from the trivial ‘fake’ decoherence by the T_2^* process, we have examined the decoherence in single-system dynamics before taking the ensemble average [4, 5]. The study of the single-system dynamics reveals a wealth of important effects [4, 5] which were missing in the density matrix cluster expansion theory [7, 8] where with inhomogeneous broadening only the spin echo signals are of interest. For example, we found that unless the external field is very strong, the off-diagonal hyperfine interaction has significant effect on the T_2 decoherence in free-induction decay (FID) even though its effect is virtually removed from the spin echo signals [4]. Thus the spin echo decay time is substantially different from the FID time with the inhomogeneous broadening effect excluded. Moreover, we discovered a recovery of the lost spin coherence by disentanglement at times away from the spin echo [5]. The concept of disentanglement led us naturally to a systematic scheme of using pulse sequences to preserve the spin coherence, of which the Carr–Purcell sequence, studied also by the Das Sarma group [8], is a special example. In this paper, we will try to give a comprehensive account of our quantum theory on the T_2 problem, taking into account both the hyperfine interaction and the intrinsic nuclear spin interaction.

Under the condition of intermediate to strong field, the Hamiltonian of the electron–nuclear spin system is reduced to the form $\hat{H} = \sum_{\pm} |\pm\rangle \hat{H}_{\pm} \langle \pm|$ which is diagonal in the electron spin eigenstate basis $\{|\pm\rangle\}$, where \hat{H}_{\pm} are the nuclear bath Hamiltonians depending on the electron spin states. The entanglement picture for the pure dephasing (T_2 decoherence) of the electron spin [43] is depicted as follows. Let the electron–nuclear spin system start from a product (i.e. unentangled) state, $(C_+|+\rangle + C_-|-\rangle) \otimes |\mathcal{J}\rangle$. The nuclear spin state $|\mathcal{J}\rangle$ would be driven by the Hamiltonians \hat{H}_{\pm} to the states $|\mathcal{J}^{\pm}(t)\rangle$ corresponding to the electron states $|\pm\rangle$, respectively. Thus the electron–nuclear spins would evolve into an entangled state, $C_+|+\rangle \otimes |\mathcal{J}^+(t)\rangle + C_-|-\rangle \otimes |\mathcal{J}^-(t)\rangle$. The off-diagonal element of the reduced density matrix of the electron spin $\rho_{+,-}^e(t) = C_-^* C_+ \langle \mathcal{J}^-(t) | \mathcal{J}^+(t) \rangle$ measures the electron spin coherence. Therefore, the bifurcated evolution of the nuclear bath leads to the loss of electron spin coherence. From the viewpoint of quantum measurement, the electron spin states are registered by different nuclear bath states and a measured quantum object sustains no coherence in the measurement basis [45, 46].

Within the timescale when the irreversible leakage of the quantum coherence of a mesoscopic bath into the macroscopic environment is negligible, it is still possible to control the mesoscopic bath dynamics [47] and hence the decoherence of the quantum object embedded in the bath [5]. As illustrated in figure 1, the bath evolution pathways in the Hilbert space corresponding to opposite electron spin states will exchange their evolution directions when the electron spin is flipped:

$$\begin{aligned} & (C_+|+\rangle + C_-|-\rangle) \otimes |\mathcal{J}\rangle \\ \longrightarrow & C_+|+\rangle \otimes |\mathcal{J}^+(t)\rangle + C_-|-\rangle \otimes |\mathcal{J}^-(t)\rangle \\ \longrightarrow & C_+|-\rangle \otimes |\mathcal{J}^+(t')\rangle + C_-|+\rangle \otimes |\mathcal{J}^-(t')\rangle. \end{aligned}$$

Thereafter, the two bath pathways $|\mathcal{J}^+(t')\rangle$ and $|\mathcal{J}^-(t')\rangle$ intersect at some later time, i.e. $|\mathcal{J}^+(t')\rangle \cong |\mathcal{J}^-(t')\rangle = |\mathcal{J}(t')\rangle$. At the intersection, the electron spin is disentangled from the bath and its lost coherence is recovered (by the controlled erasure of the quantum information registered in the bath). Such a discovery of the reversal of coherent dynamics can be traced back to the early theoretical and experimental findings on the Loschmidt echo in NMR [48, 49].




Figure 1. Schematic illustration of the bifurcation of the bath state evolution conditioned on the electron spin up or down state and the exchange of the evolution direction of the bath pathways when the electron spin is flipped.

The recovery of the lost coherence (recoherence) by quantum disentanglement is fundamentally different from the conventional spin echo realized by the refocusing of the random phase in an inhomogeneously broadened ensemble. In general, the disentanglement can occur at a time different from the spin echo time. For example, by a single flip of the electron spin at $t = \tau$, there will be a prominent recoherence at a magic time $\sqrt{2}\tau$ as a consequence of disentanglement [5] while the conventional spin echo would occur at 2τ . When the electron spin is observed in ensemble measurement (realized either by using many similar QDs or by cycling measurements on a single dot), the disentanglement-induced recoherence will be concealed by the inhomogeneous broadening unless it is forced to take place at a spin echo time by proper design of pulse sequences.

The recoherence by disentanglement can be exploited as a coherence protection scheme in lieu of the dynamical decoupling schemes developed for NMR spectroscopies [48, 50, 51] and recently for quantum computation [52]–[57]. While the dynamical decoupling schemes seek to eliminate the object–bath interaction through rapid rotation of a quantum object, the disentanglement scheme focuses on controlling the wavefunction evolution of the bath and in general does not lead to a vanishing object–bath coupling. It will be shown that in the disentanglement scheme, the controlling small parameter for coherence protection is determined by the interactions within the bath instead of the object–bath coupling, in contrast with the dynamical decoupling schemes.

This paper is organized as follows: section 2 describes a specific model system (a self-assembled InAs QD) and summarizes the theoretical ingredients for solving the electron–nuclear spin dynamics. The FID in single-system dynamics (without ensemble average) and the Hahn echo in ensemble dynamics are studied in section 3 and 4, respectively. Section 5 presents a close examination of the single-system dynamics in the single-pulse Hahn-echo configuration, revealing a coherence recovery due to disentanglement at a magic time which would otherwise be invisible in ensemble-averaged signals. Section 6 studies the disentanglement under control of pulse sequences, in particular, the concatenated sequences. Further discussions including the comparison between the quantum disentanglement and the dynamical decoupling are presented in the summary section. Some technical details are given in the appendices.

2. Theory

2.1. The model

The system consists of an electron with spin-1/2 \hat{S}_e and N nuclear spins, $\hat{J}_{n,\alpha}$, with Zeeman energies Ω_e and $\omega_{n,\alpha}$ under a magnetic field B_{ext} , respectively, where n denotes the position and α denotes the isotope type. Hereafter the subscript α will be absorbed into n unless it is needed for clarity. InAs has the zincblende structure, with In and As ions located in two interpenetrating face-centred cubic lattices. The natural isotope abundance in InAs materials is 100, 4.28 and 95.72% for ^{75}As , ^{113}In and ^{115}In , respectively. All the isotopes have nonzero nuclear spin moments, namely, $J_{^{75}\text{As}} = 3/2$ and $J_{^{113}\text{In}} = J_{^{115}\text{In}} = 9/2$. The self-assembled InAs QD under typical growth condition is modelled as a rectangular quantum box with the growth direction along [001] and the in-plane extension directions [110] and [1 $\bar{1}$ 0] [58]. The electron is assumed to be confined by hardwall potential and the envelope wavefunction of the ground state is

$$f(\mathbf{r}) = \prod_{\mathbf{a}} \sqrt{\frac{2}{L_{\mathbf{a}}}} \cos \frac{\pi r_{\mathbf{a}}}{L_{\mathbf{a}}} \theta(L_{\mathbf{a}} - 2|r_{\mathbf{a}}|), \quad (1)$$

where $r_{\mathbf{a}}$ is the coordinate in the \mathbf{a} direction ([110], [1 $\bar{1}$ 0], or [001]), and $L_{\mathbf{a}}$ is the dimension of the dot along the indicated direction. The model system is illustrated in figure 2.

The first-principles Hamiltonian for the electron–nuclear spin system includes the electron–nuclear hyperfine interaction and various intrinsic nuclear–nuclear interactions. A detailed discussion was given in [4]. Under a moderate magnetic field ($B_{\text{ext}} > 0.1$ T), the electron spin flip is virtually suppressed due to the large Zeeman energy mismatch between the electron and nuclei. Virtual flip–flops of the electron spin, however, could mediate an extrinsic interaction between nuclear spins even when they are well separated in space (as illustrated in figure 3) [4]. Furthermore, the non-secular part of the nuclear spin interaction which does not conserve the total Zeeman energy, including flip–flops of hetero-nuclear pairs and single-spin flips, are neglected. Thus the total effective Hamiltonian is reduced to the form

$$\hat{H} = \hat{H}_e + \hat{H}_N + \sum_{\pm} |\pm\rangle \hat{H}_{\pm} \langle \pm|, \quad (2)$$

in the limit of long longitudinal relaxation time ($T_1 \rightarrow \infty$), where $|\pm\rangle$ are the eigenstates of \hat{S}_e^z , $\hat{H}_e \equiv \Omega_e \hat{S}_e^z$, $\hat{H}_N \equiv \omega_n \hat{J}_n^z$, and the nuclear spin interaction

$$\hat{H}_{\pm} = \pm \hat{H}_A + \hat{H}_B + \hat{H}_D \pm \hat{H}_E, \quad (3)$$

with

$$\hat{H}_A \equiv \sum_{n \neq m}' \frac{a_n a_m}{4\Omega_e} \hat{J}_n^+ \hat{J}_m^- \equiv \sum_{n \neq m}' A_{n,m} \hat{J}_n^+ \hat{J}_m^-, \quad (4a)$$

$$\hat{H}_B \equiv \sum_{n \neq m}' B_{n,m} \hat{J}_n^+ \hat{J}_m^-, \quad (4b)$$




Figure 2. (a) Schematics of a quantum box model for an InAs QD. (b) Schematics of an electron (the shadow) and one layer of nuclear spins in the QD. The two boxes in dotted lines indicate two possible choices of boundary of the nuclear spin bath, which are relatively arbitrary due to the interaction between nuclei within and without the boundary. When the hyperfine interaction dominates over the nuclear spin interaction, such arbitrariness has negligible effects on calculation of the electron spin decoherence as long as all the nuclei in direct contact with the electron spin have been enclosed.




Figure 3. Hyperfine-mediated interaction between two distant nuclear spins via the electron (top line).

$$\hat{H}_D \equiv \sum_{n < m} D_{n,m} \hat{J}_n^z \hat{J}_m^z, \quad (4c)$$

$$\hat{H}_E \equiv \sum_n (a_n/2) \hat{J}_n^z \equiv \sum_n E_n \hat{J}_n^z. \quad (4d)$$

Table 1. InAs material parameters, where γ_α is the nuclear gyromagnetic ratios, $\gamma_\alpha^{\text{ex}}$ the effective nuclear gyromagnetic ratios for the pseudo-exchange interaction, γ_e the electron gyromagnetic ratio, γ_e^* the effective electron gyromagnetic ratio in the QD, and \mathcal{A}_α the hyperfine constants.

	⁷⁵ As	¹¹³ In	¹¹⁵ In
Abundance	100%	4.28%	95.72%
Spin moment J_α	3/2	9/2	9/2
γ_α ($10^6 \text{ s}^{-1} \text{ T}^{-1}$)	+45.8	+58.5	+58.6
$\gamma_\alpha^{\text{ex}}$ ($10^6 \text{ s}^{-1} \text{ T}^{-1}$) ^a	+34.0	+70.1	+70.2
\mathcal{A}_α (10^9 s^{-1}) ^b	+69.8	+85.1	+85.3
$\gamma_e = +0.176$, $\gamma_e^* = -0.132$ ($10^{12} \text{ s}^{-1} \text{ T}^{-1}$) ^c			
Lattice constant $c_0 = 6.06 \text{ \AA}$			

^aSee details in [4, 59, 60].

^bEstimated with the method in [61].

^c g -factor taken to be -1.5 .

Table 2. The characteristic energy scales in an InAs QD with dimensions $35 \times 35 \times 6 \text{ nm}^3$ ($N \sim 0.3 \times 10^6$), under a field of $B_{\text{ext}} \sim 10$ tesla.

Units ^a	Ω_e	ω_α	E_n	$B_{n,m}, D_{n,m}$	$A_{n,m}$
10^6 s^{-1}	10^6	500	1	10^{-4}	10^{-6}
μeV	10^3	0.5	10^{-3}	10^{-7}	10^{-9}
mK	10^4	5	10^{-2}	10^{-6}	10^{-8}

^aThroughout this paper, the units are chosen such that the Plank constant \hbar and the Boltzmann constant k_B are unity.

Here the summation with a prime runs over only the homo-nuclear pairs, \hat{H}_A denotes the extrinsic interaction mediated by the hyperfine interaction, \hat{H}_B denotes the off-diagonal part of the intrinsic nuclear interaction and \hat{H}_D the diagonal part, and \hat{H}_E is the diagonal part of the contact hyperfine interaction with amplitude $a_n = \mathcal{A}_\alpha c_0^3 |f(\mathbf{R}_n)|^2$. The material parameters are given in table 1 and the typical energy scales are estimated in table 2. The hyperfine interaction has a typical energy scale $E_n \sim a_n \sim 10^6 \text{ s}^{-1}$ for a dot with about 10^6 nuclei [61]. The intrinsic nuclear spin–spin interaction is effectively finite-ranged with the near-neighbour coupling $B_{n,m} \sim D_{n,m} \sim 10^2 \text{ s}^{-1}$. The extrinsic interaction is ‘infinite-ranged’ (coupling any two nuclear spins within the QD) and has opposite signs for opposite electron spin states. The extrinsic interaction, in contrast with the intrinsic one, depends on the external field, and has an energy scale $A_{n,m} \sim 1$ to 10 s^{-1} for field strength B_{ext} varying from 10 to 1 tesla.

We reiterate that our model has included both the extrinsic hyperfine-mediated and the intrinsic nuclear–nuclear spin interactions. On one hand, without the extrinsic interaction included [7]–[9], the FID in single system dynamics would not be correctly described, especially for relatively weak field. On the other hand, without the intrinsic interaction considered [17]–[22], the theory could not account for the FID in the strong field case when the hyperfine-mediated interaction is suppressed [4, 21, 22, 24], nor for spin echo signals where the hyperfine interaction effect is removed in the leading order [4, 23].

2.2. General formalism

The electron–nuclear spin system is assumed initially prepared in a product state⁴

$$\hat{\rho}(0) = \hat{\rho}^e(0) \otimes \hat{\rho}^N. \quad (5)$$

where the nuclear spins are in a thermal state with temperature T . The electron spin can start from a pure state or a mixed state⁵. The state evolution is determined by the Louville equation

$$\partial_t \hat{\rho}(t) = -i[\hat{H}, \hat{\rho}(t)]. \quad (6)$$

The reduced density matrix of the electron spin is obtained by partial trace over the nuclear spins as

$$\hat{\rho}^e(t) = \text{Tr}_N \hat{\rho}(t), \quad (7)$$

which is related to the initial state through the correlation superoperator $\hat{\mathcal{L}}$, namely

$$\rho_{\mu,v}^e(t) = \sum_{\mu',v'} \mathcal{L}_{\mu,v;\mu',v'}(t) \rho_{\mu',v'}^e(0), \quad (8)$$

where $\rho_{\mu,v}^e \equiv \langle \mu | \hat{\rho}^e | v \rangle$, and $|\mu\rangle, |v\rangle \in \{|+\rangle, |-\rangle\}$. The electron spin relaxation is quantified by the correlation function $\mathcal{L}_{\mu,v;\mu',v'}$.

Due to the block diagonal form of the reduced Hamiltonian equation (2), the correlation function has following properties

$$\mathcal{L}_{\mu,v;\mu',v'}(t) = \mathcal{L}_{\mu,v}(t) \delta_{\mu,\mu'} \delta_{v,v'}, \quad (9a)$$

$$\mathcal{L}_{\mu,\mu}(t) = 1, \quad (9b)$$

$$\begin{aligned} \mathcal{L}_{+,-}(t) &= \mathcal{L}_{-,+}^*(t) \\ &= e^{-i\Omega_e t} \text{Tr}_N [\hat{\rho}^N e^{+i\hat{H}_- t} e^{-i\hat{H}_+ t}], \end{aligned} \quad (9c)$$

where the last equation has been expressed for the FID and can be straightforwardly extended to the dynamics under pulse control. No longitudinal relaxation remains ($T_1 = \infty$) after the elimination of the electron spin flip processes by the magnetic field. Since the Zeeman energy \hat{H}_0 results in only a global phase-factor $e^{-i\Omega_e t}$, from now on we will consider only the interaction Hamiltonian $\hat{H}_1 \equiv \hat{H} - \hat{H}_0$ by dropping the trivial phase-factor.

Since the nuclear Zeeman energy under the external field dominates over the hyperfine interaction and the nuclear spin interaction and the temperature to be considered is much higher than the interaction energy, the thermal nuclear spin state can be taken as

$$\hat{\rho}^N \cong e^{-\hat{H}_N/T} = \sum_{\mathcal{J}} P_{\mathcal{J}} |\mathcal{J}\rangle \langle \mathcal{J}|, \quad (10)$$

⁴ It is also interesting to consider the electron spin decoherence and the nuclear spin dynamics for a system with certain initial electron–nuclear entanglement, which will be covered in our future publications.

⁵ The electron spin being in a mixed state means that it is entangled with its ‘environment’. Since we have assumed the electron and the nuclei are initially in a product state, the mixedness must come from other sources, such as the photon noise in optical pump and control of the electron spin.

where $|\mathcal{J}\rangle \equiv \bigotimes_n |j_n\rangle$ is an eigenstate of \hat{H}_N and $P_{\mathcal{J}} = \prod_n p_{j_n}$ is the probability distribution with

$$p_{j_n,\alpha} \equiv \frac{e^{-j_n,\alpha\omega_n/T}}{\sum_{j=-J_\alpha}^{+J_\alpha} e^{-j\omega_n/T}}, \quad (11)$$

giving the population of the single-spin state $|j_n\rangle$. We note that such a nuclear spin state has no off-diagonal coherence (between the Zeeman energy eigenstates). The correlation function of the electron spin coherence, expressed in terms of the wavefunction overlap of the nuclear spins, is

$$\mathcal{L}_{+,-}(t) = \sum_{\mathcal{J}} P_{\mathcal{J}} \langle \mathcal{J}^-(t) | \mathcal{J}^+(t) \rangle, \quad (12)$$

with $|\mathcal{J}^\pm(t)\rangle \equiv \exp(-i\hat{H}_\pm t) |\mathcal{J}\rangle$. The correlation function $\mathcal{L}_{+,-}(t)$ is closely related to the Loschmidt echo in [63]–[65]. Being independent of the electron spin initial state, $\mathcal{L}_{+,-}(t)$ will be equated with the electron spin coherence without causing confusion. The decoherence by entanglement is transparent from equation (12): with the nuclear spin states $|\mathcal{J}^\pm(t)\rangle$ driven by different Hamiltonians \hat{H}_\pm , the electron–nuclear spin state $C_+|+\rangle \otimes |\mathcal{J}^+(t)\rangle + C_-|-\rangle \otimes |\mathcal{J}^-(t)\rangle$ becomes an entangled one when $\langle \mathcal{J}^-(t) | \mathcal{J}^+(t) \rangle < 1$, so the electron spin coherence is lost.

2.3. Essential assumptions

The method to be used in this paper consists of two steps. Firstly, the electron spin decoherence in *ensemble dynamics* is factorized into two parts: the dephasing due to static inhomogeneous broadening, and the decoherence due to the dynamical entanglement between the electron and the nuclei in *single-system dynamics*. Secondly, the nuclear spin dynamics is solved with the pair-correlation approximation. The following assumptions are essential for the theory.

1. *Mesoscopia*: firstly, the QD should be small enough to guarantee the dominance of the hyperfine interaction over the nuclear spin interaction so that the isolation of a mesoscopic bath is possible. Secondly, the bath size (i.e. the number of nuclear spins in the QD, N) should be large enough to have genuine decoherence (i.e. the Poincaré time is much greater than the spin relaxation time T_1 and T_2). Thirdly, $N \gg \sqrt{N}$ is required so that the central limit theorem in statistics can be utilized for the factorization of the single-system dynamics from the inhomogeneous broadening effect. And finally, to justify the pair-correlation approximation, the bath size N is bounded so that the number of nuclear pair-flips in the relevant timescale is small compared with N (see appendix B).
2. *Moderate to strong field*: the electron Zeeman energy Ω_e be much greater than the hyperfine energy a_n so that the electron spin flip is suppressed [4, 17, 21].⁶
3. *Finite temperature*: the temperature be higher than or comparable to the nuclear Zeeman energy ($T \gtrsim \omega_\alpha \sim 1$ mk). Otherwise, in a near fully polarized nuclear spin bath under an extremely low-temperature, the pair flip–flops occur only among a few available nuclear spins and hence are highly correlated, in which the pair-correlation approximation is

⁶ In low fields, it may be possible to treat the off-diagonal hyperfine interaction as elementary excitations within the pair-correlation approximation as well as the nuclear spin pair-flips are. This generalization, however, is beyond the scope of this paper.

invalid. Also, the temperature should be low enough to prevent the phonon scattering effect ($T \lesssim 1$ K) [27]–[30].

4. *Short-time dynamics*: the timescales under consideration should be short compared with the inverse nuclear spin interaction strength (~ 1 ms) so that the number of pair-flips over time is small compared with N . The timescales of interest in the decoherence problem include the coherence memory time, pulse delay time in control sequences, and the total duration of a control pulse sequence.

The other approximations in the model, including the assumptions on the QD shape, the electron wavefunction, and the specific form of the spin interactions, are inconsequential to the theory.

2.4. Ensemble and single-system dynamics

The electron spin decoherence given in equation (9) includes two contributions, namely, the thermal fluctuations due to the ensemble average of different nuclear spin configurations $\{|\mathcal{J}\rangle\langle\mathcal{J}|\}$, and the quantum entanglement due to the dynamical evolution starting from a pure state $|\mathcal{J}\rangle$, which is conditioned on the electron spin state. We use the term *single-system dynamics* to denote the quantum evolution governed by the Schrödinger equation

$$i\partial_t|\mathcal{J}^\pm(t)\rangle = H_\pm|\mathcal{J}^\pm(t)\rangle, \quad (13)$$

and *ensemble dynamics* to denote the ensemble average of the single-system dynamics.

The ensemble dynamics was studied in the formalism of density matrix with the pair-correlation approximation [7]. Many important features of the quantum entanglement process, however, could have been overlooked without resolving the single-system dynamics since the static inhomogeneous broadening usually has much stronger effects on the decoherence than the dynamical entanglement. We choose a different procedure to study the problem. Namely, we will first solve the single-system dynamics starting from a certain nuclear spin configuration, and then construct the ensemble dynamics via

$$\mathcal{L}_{+,-}^{\mathcal{J}}(t) \equiv \langle\mathcal{J}^-(t)|\mathcal{J}^+(t)\rangle, \quad (14a)$$

$$\mathcal{L}_{+,-}(t) = \sum_{\mathcal{J}} P_{\mathcal{J}} \mathcal{L}_{+,-}^{\mathcal{J}}(t). \quad (14b)$$

Such a construction of the ensemble dynamics, in general, would require sampling a large number of initial states $|\mathcal{J}\rangle$ in the thermal ensemble.

When the system is sufficiently large and the temperature is appreciable for the nuclear spins, the decoherence due to quantum fluctuations in single-system dynamics (equation 14a) is almost the same for all possible initial nuclear configurations in the thermal ensemble, except for a global phase factor related to the static Overhauser field resulting from the diagonal hyperfine interaction. This is confirmed by numerical verification (see appendix A). Here we give a physical argument: for a sufficiently large number of nuclear spins far from being fully polarized, the number of spins available for pair-wise flip–flops is large, so the electron spin decoherence is virtually determined by the nuclear spin excitation spectra (or density of states). By the statistical central limit theorem, the excitation spectrum of a large system is the same (up to a relative variance $\sim 1/\sqrt{N}$) for different initial states $|\mathcal{J}\rangle$. Then the electron spin decoherence can be

factorized into a contribution from the static inhomogeneous broadening \mathcal{L}^{inh} and one from the quantum fluctuation in single-system dynamics \mathcal{L}^{s} , defined as

$$\mathcal{L}_{+,-}(t) \cong \mathcal{L}_{+,-}^{\text{inh}}(t)\mathcal{L}_{+,-}^{\text{s}}(t), \quad (15a)$$

$$\mathcal{L}_{+,-}^{\text{s}}(t) \equiv |\langle \mathcal{J}^-(t) | \mathcal{J}^+(t) \rangle|, \quad (15b)$$

$$\mathcal{L}_{+,-}^{\text{inh}}(t) = \sum_{\mathcal{J}} P_{\mathcal{J}} e^{-iE_{\mathcal{J}}t} = \int P(E) e^{-iEt} dE, \quad (15c)$$

where the hyperfine energy due to the local Overhauser field in a certain nuclear spin configuration is

$$E_{\mathcal{J}} \equiv \sum_n j_n a_n, \quad (16)$$

and the inhomogeneous broadening distribution is defined as

$$P(E) \equiv \sum_{\mathcal{J}} P_{\mathcal{J}} \delta(E - E_{\mathcal{J}}). \quad (17)$$

The quantum fluctuation effect in the single-system dynamics can be evaluated for an initial configuration $|\mathcal{J}\rangle$ randomly selected from the ensemble, with the global phase factor absorbed into the inhomogeneous broadening.

The effect of the static inhomogeneous broadening can be calculated by neglecting the nuclear spin interaction (setting $\hat{H}_A = \hat{H}_B = 0$). Without the flip-flop dynamics, the nuclear spin ensemble would be a random distribution of frozen configurations, which, as a standard multinomial distribution for a large system, leads to a Gaussian distribution of the Overhauser field

$$P(E_{\mathcal{J}}) = \frac{1}{\sqrt{2\pi}\Gamma_2^*} e^{-(E_{\mathcal{J}}-E_0)^2/(2\Gamma_2^*)^2}, \quad (18)$$

where the averaged local field E_0 and the inhomogeneous broadening Γ_2^* are

$$E_0 \cong \sum_{\alpha} (2N_{\alpha} A_{\alpha} N^{-1} \bar{j}_{\alpha}) \equiv \sum_{\alpha} E_{0,\alpha}, \quad (19a)$$

$$\Gamma_2^* \cong \sqrt{\sum_{\alpha} (4N_{\alpha} A_{\alpha}^2 N^{-2} \bar{j}_{\alpha}^2 - E_{0,\alpha}^2 N_{\alpha}^{-1})}, \quad (19b)$$

respectively, with $\bar{j}_{\alpha} \equiv \sum_j j p_{\alpha,j}$, $\bar{j}_{\alpha}^2 \equiv \sum_j j^2 p_{\alpha,j}$, and N_{α} denoting the number of α -type nuclei ($N \equiv \sum_{\alpha} N_{\alpha}$). Thus we reproduce the expressions for the T_2^* dephasing obtained in [17]. A typical example of the Overhauser field distribution is shown in figure 4. The electron spin dephasing by inhomogeneous broadening is a Gaussian decay

$$\mathcal{L}_{+,-}^{\text{inh}}(t) = e^{-iE_0 t - (t/T_2^*)^2}, \quad (20)$$

with the effective dephasing time $T_2^* \equiv \sqrt{2}/\Gamma_2^*$. The inhomogeneous broadening effect can be removed by spin echo. In the rotating frame rested on the electron spin, the Overhauser field changes its sign each time the electron spin is flipped by a short pulse. Thus under the flipping




Figure 4. Distribution of the Overhauser field for a thermal ensemble of nuclear spins in an InAs QD of size $34 \times 34 \times 3 \text{ nm}^3$ at a temperature of 1 K and under an external field of 1 tesla. The dots in the figure are obtained by numerical simulation of 40 000 random samples, and the curve is calculated with equations (18)–(19). The arrows indicate four randomly selected nuclear spin configurations to be used for figure A.1 in appendix A.

operation of a sequence of pulses applied at $t = t_1, t_2, \dots, t_n$, the dephasing by the inhomogeneous broadening becomes

$$\mathcal{L}_{+,-}^{\text{inh}}(t) = \sum_{\mathcal{J}} P_{\mathcal{J}} e^{-iE_{\mathcal{J}}\bar{t}} = e^{-iE_0\bar{t} - \bar{t}^2/(T_2^*)^2}, \quad (21)$$

with $\bar{t} \equiv t_1 - (t_2 - t_1) + (t_3 - t_2) \cdots - (-1)^n(t - t_n)$. Obviously, the phase accumulation from the random Overhauser field is cancelled at $\bar{t} = 0$, leading to the spin echo. At the spin-echo time, the electron spin decoherence results solely from the dynamical quantum entanglement.

Under typical conditions considered in this paper, T_2^* is found to be of the order of nanoseconds, consistent with various experimental measurements [37, 39, 40, 65]. As will be shown below, in agreement with the available experimental data [40, 41, 66], the electron spin decoherence by quantum fluctuations in single-system dynamics has a timescale in the order of microseconds. Thus, from equation (15), the ensemble dynamics will be dominated by inhomogeneous broadening and the single-system dynamics is virtually invisible in ensemble experiments, unless the inhomogeneous broadening effect is removed by spin echo. The decay of spin-echo signals is usually attributed to the pure decoherence due to quantum fluctuations. We have shown, however, by examining directly the single-system dynamics, the flipping pulses have nontrivial effects on the nuclear spin dynamics and, therefore, on the dynamical entanglement, making the spin-echo decay time significantly different from the FID time in single-system dynamics [4]. Such a difference would not be realized without resolving the single-system dynamics from the ensemble average [7]–[9].

From now on, we will concentrate on the single-system dynamics, and show a few surprising effects which would otherwise be concealed by inhomogeneous broadening in ensemble dynamics. It should be emphasized that the single-system dynamics in a QD at a finite temperature is not a mathematical idealization but has measurable effects. For example, a projective measurement of the local Overhauser field could be used to limit the nuclear spin configurations by post-selection and thus to observe single-electron dynamics without spin echo [20], [67]–[69]. The single-system dynamics is basic to quantum technologies such as quantum computation which cannot be performed in ensembles for scalability to large systems without exponential explosion in resource [70].

2.5. Pair-correlation approximation and pseudo-spin model for nuclear spin dynamics

With the Hamiltonian given in equation (4), the quantum evolution of the nuclear spins is driven by the pair-wise homo-nuclear spin flip–flops as the elementary excitations. The transition for the pair-flip driven by the operator $J_n^+ J_m^-$ is

$$|j_n\rangle_n |j_m\rangle_m \longrightarrow |j_n + 1\rangle_n |j_m - 1\rangle_m. \quad (22)$$

The transition between the electron–nuclear spin many-body states through the pair-flip is denoted by

$$|\pm\rangle|\mathcal{J}\rangle \longrightarrow |\pm\rangle|\mathcal{J}, k\rangle, \quad (23)$$

where k is shorthand for the pair-flip in equation (22). The transition described in equation (23) is characterized by the matrix elements $\pm A_k + B_k$ and the energy costs $D_k \pm E_k$, which are derived from the microscopic models as follows

$$A_k \equiv \langle \mathcal{J}, k | \hat{H}_A | \mathcal{J} \rangle = \frac{a_n a_m}{4\Omega_e} \sqrt{J_n(J_n + 1) - j_n(j_n + 1)} \sqrt{J_m(J_m + 1) - j_m(j_m - 1)}, \quad (24a)$$

$$B_k \equiv \langle \mathcal{J}, k | \hat{H}_B | \mathcal{J} \rangle = B_{n,m} \sqrt{J_n(J_n + 1) - j_n(j_n + 1)} \sqrt{J_m(J_m + 1) - j_m(j_m - 1)}, \quad (24b)$$

$$\begin{aligned} D_k &\equiv \langle \mathcal{J}, k | \hat{H}_D | \mathcal{J}, k \rangle - \langle \mathcal{J} | \hat{H}_D | \mathcal{J} \rangle \\ &= \sum_{n'} D_{n,n'} j_{n'} - \sum_{m'} D_{m,m'} j_{m'} - D_{n,m} + D_{n,n} (j_n + 1) - D_{m,m} (j_m - 1), \end{aligned} \quad (24c)$$

$$E_k \equiv \langle \mathcal{J}, k | \hat{H}_E | \mathcal{J}, k \rangle - \langle \mathcal{J} | \hat{H}_E | \mathcal{J} \rangle = (a_n - a_m)/2. \quad (24d)$$

The basic processes for the electron spin decoherence may be described as follows. The off-diagonal part of the nuclear spin interaction (including the intrinsic one \hat{H}_B and the extrinsic hyperfine-mediated one \hat{H}_A) causes the pair-wise nuclear spin flip–flops, leading to a fluctuating local Overhauser field and in turn a random dynamical phase for the electron spin. In the quantum picture, the entanglement is developed because the quantum evolution of the nuclear spins driven by the Hamiltonians $\pm \hat{H}_A + \hat{H}_B + \hat{H}_D \pm \hat{H}_E$ is conditioned on the electron spin state through the \pm signs originating from the hyperfine interaction.

As discussed in [4] and [7] and further checked for pulse sequence control cases in appendix C, within a time t much smaller than the inverse nuclear interaction strength, the total number of pair-flip excitations N_{flip} is much smaller than the number of nuclei N . The probability of having pair-flips correlated is estimated to be $P_{\text{corr}} \sim 1 - e^{-qN_{\text{flip}}^2/N}$ (q being the number of nearest neighbours) [7], which, as also shown by *a posteriori* numerical check (see appendix C), is bounded by $\sim 10\%$ in the worst scenario studied in this paper. Thus, the pair-flips as elementary excitations from the initial state can be treated as independent of each other, with a relative error $\epsilon \lesssim P_{\text{corr}}$ (see appendix D). Then the single-system dynamics $|\mathcal{J}^\pm(t)\rangle$ can be described by the excitation of pair-correlations as non-interacting quasi-particles from the ‘vacuum’ state $|\mathcal{J}\rangle$, driven by the ‘low-energy’ effective Hamiltonian,

$$\hat{H}_{\mathcal{J}}^\pm = \sum_k \hat{\chi}_k^\pm \equiv \sum_k \chi_k^\pm \cdot \hat{\sigma}_k / 2, \quad (25)$$

which has been written in such a way that each pair-flip is treated as a pseudo-spin $1/2$, represented by the Pauli matrix $\hat{\sigma}_k$, with k labelling all possible pair-flips. The time evolution from the initial state $|\mathcal{J}\rangle$ can be viewed as the rotation of the pseudo-spins, initially all polarized along the $-z$ pseudo-axis: $\bigotimes_k |\downarrow\rangle_k$, under the effective pseudo-magnetic field,

$$\chi_k^\pm \equiv (\pm 2A_k + 2B_k, 0, D_k \pm E_k), \quad (26)$$

for the electron spin state $|\pm\rangle$, respectively. Such a treatment of the nuclear spin correlations amounts to taking into account all the pairwise correlations and neglecting the higher-orders. The pair-correlation approximation, equivalent to the lowest order cluster expansion in [7], corresponds to the leading term in the linked cluster expansion [6] and the density matrix cluster expansion [9].

The entanglement between the electron spin and the pseudo-spins, and hence the electron spin decoherence, are developed as the pseudo-spins precess about different pseudo-fields χ_k^\pm corresponding to different electron spin states $|\pm\rangle$. Namely, the product state

$$(C_+|+\rangle + C_-|-\rangle) \bigotimes_k |\downarrow\rangle_k, \quad (27)$$

will evolve into an entangled one

$$C_+|+\rangle \bigotimes_k |\psi_k^+(t)\rangle + C_-|-\rangle \bigotimes_k |\psi_k^-(t)\rangle, \quad (28)$$

with

$$|\psi_k^\pm(t)\rangle = \exp(-i\hat{\chi}_k^\pm t) |\downarrow\rangle = e^{-i\chi_k^\pm \cdot \hat{\sigma}_k t/2} |\downarrow\rangle. \quad (29)$$

The electron spin coherence is determined by the overlap between the ‘conjugate’ states $|\psi_k^+(t)\rangle$ and $|\psi_k^-(t)\rangle$ of the pseudo-spins,

$$\mathcal{L}_{+,-}^s(t) = \prod_k |\langle \psi_k^-(t) | \psi_k^+(t) \rangle|. \quad (30)$$

The pseudo-spin states have a geometrical representation as Bloch vectors,

$$\sigma_k^\pm \equiv \langle \psi_k^\pm | \hat{\sigma}_k | \psi_k^\pm \rangle / 2. \quad (31)$$

Figure 5 shows the evolution of two conjugate pseudo-spin states $|\psi_k^\pm(t)\rangle$ in two paths of the conjugate Bloch vectors σ_k^\pm on a sphere. The overlap between the conjugate pseudo-spin states is related to the distance between the two vectors $\delta_k = |\sigma_k^- - \sigma_k^+|$ by

$$|\langle \psi_k^-(t) | \psi_k^+(t) \rangle|^2 = 1 - \delta_k^2. \quad (32)$$

Thus, the square of the distance δ_k^2 is the distinguishability between the two conjugate states. This geometrical picture gives an interpretation of the decoherence process as a measurement. The ‘measuring device’ composed of nuclear spins evolves in two pathways, each for an electron spin eigenstate, up or down. When the sum of the distinguishability between the conjugate states of all pair-flips is large enough, the ‘device’ states are ‘macroscopically’ distinguishable. The coherence between the basis states is destroyed by the measurement [46]. The pseudo-spin picture will provide us with a physical guide to the decoherence control by pulse sequences, which is not available from the density matrix expansion method [7].




Figure 5. The Bloch vectors for the pseudo-spin states $|\psi_k^\pm(t)\rangle$ precess about the pseudo-fields χ_k^\pm for the electron spin states $|\pm\rangle$, respectively.

3. Free-induction decay

The separation of the single-system dynamics from the ensemble dynamics offers an opportunity to study the FID of electron spin coherence due to dynamical entanglement, which would otherwise be concealed by the much stronger effect of inhomogeneous broadening and could not be resolved in ensemble dynamics study [7].

The pseudo-spin evolution is directly calculated from equation (29) to be

$$|\psi_k^\pm(t)\rangle = \left(\cos \frac{\chi_k^\pm t}{2} - i \hat{\sigma}_k \cdot \sin \frac{\chi_k^\pm t}{2} \right) | \downarrow \rangle, \quad (33)$$

where the sine function of a vector \mathbf{v} is defined as $\sin(\mathbf{v}) \equiv (\mathbf{v}/v) \sin(v)$. The distinguishability between the conjugate pseudo-spin states $|\psi_k^\pm\rangle$ is

$$\begin{aligned} \delta_k^2 = & t^4 (B_k E_k - A_k D_k)^2 \operatorname{sinc}^2 \frac{\chi_k^+ t}{2} \operatorname{sinc}^2 \frac{\chi_k^- t}{2} \\ & + t^2 \left[(B_k + A_k) \cos \frac{\chi_k^- t}{2} \operatorname{sinc} \frac{\chi_k^+ t}{2} - (B_k - A_k) \cos \frac{\chi_k^+ t}{2} \operatorname{sinc} \frac{\chi_k^- t}{2} \right]^2. \end{aligned} \quad (34)$$

This expression is used in the numerical evaluation of the electron spin coherence.

To gain some insight into the FID features, we make two approximations which are well justified. First, the pseudo-spins are separated into two groups, \mathbb{G}_A and \mathbb{G}_B , corresponding to nonlocal and local flip-flops, respectively. For the local pair-flips ($k \in \mathbb{G}_B$), the transition matrix element is dominated by the intrinsic nuclear spin interaction as $B_k \gg A_k$ under a moderate magnetic field, while for the nonlocal pair-flips ($k \in \mathbb{G}_A$), the extrinsic hyperfine-mediated interaction dominates⁷. Thus the coherence is factorized as

$$\mathcal{L}_{+,-}^s = \mathcal{L}_{+,-}^A \times \mathcal{L}_{+,-}^B, \quad (35a)$$

⁷ The boundary between the two groups is not exact but this is not important since the pair-flips around the boundary are much weaker in the strength of the transition matrix elements than those in \mathbb{G}_B and are much fewer in number than in \mathbb{G}_A .




Figure 6. (a) The conjugate Bloch vectors for a pair-flip by the intrinsic nuclear interaction with the diagonal part neglected ($\chi_k^\pm = (B_k, 0, \pm E_k)$). (b) The same as (a) but for a nonlocal pair-flip [$\chi_k^\pm = \pm(A_k, 0, E_k)$]. (c) and (d) are the projections of (a) and (b) to the x - y -plane, respectively, in which the dotted trajectories show the effect of the diagonal nuclear interaction (D_k) (with the deviation from the solid curves exaggerated for visibility).

$$\begin{aligned} \mathcal{L}_{+,-}^{A/B} &= \prod_{k \in \mathbb{G}_{A/B}} |\langle \psi_k^-(t) | \psi_k^+ \rangle| \\ &\cong \exp \left(-\frac{1}{2} \sum_{k \in \mathbb{G}_{A/B}} \delta_k^2 \right). \end{aligned} \quad (35b)$$

The pseudo-spins in \mathbb{G}_A and those in \mathbb{G}_B have dramatically different precession behaviours. The conjugate pseudo-spins for a nonlocal pair-flip precess into opposite directions (see figures 6(b) and (d)), since both the transition amplitude (A_k) and the dominating part of energy cost (E_k) are associated with opposite signs for opposite electron spin states because of their hyperfine interaction origins. On the other hand, the intrinsic nuclear spin interaction is independent of the electron spin state, and the corresponding pseudo-spin bifurcates tangentially (see figures 6(a) and (c)). In the second approximation, we neglect the diagonal nuclear spin interaction D_k , which is justified from equation (34) with the condition $D_k \ll E_k$. This simplification can also be understood with the geometrical picture shown in figure 6. A small change of the energy cost by the diagonal nuclear interaction induces only a slight modification of the pseudo-spin precession pathways, and hence a negligible change of the distance δ_k . We will see, however, the diagonal nuclear spin interactions are important for the decoherence dynamics under control by pulse sequences.

With the two simplifications and the conditions $E_k \gg B_k, A_k$, the distinguishability between conjugate pseudo-spin states is

$$\delta_k^2 = \begin{cases} 4t^2 A_k^2 \cos^2 \frac{E_k t}{2} \text{sinc}^2 \frac{E_k t}{2} \approx 4t^2 A_k^2 & (k \in \mathbb{G}_A), \\ t^4 E_k^2 B_k^2 \text{sinc}^4 \frac{E_k t}{2} \approx t^4 E_k^2 B_k^2 & (k \in \mathbb{G}_B), \end{cases} \quad (36)$$

where the short-time approximation holds for $t \ll E_k^{-1}$. Within timescales of interest ($t \ll B_k^{-1}, A_k^{-1}$), it is always satisfied that $\delta_k^2 \ll 1$, so the coherence is approximated as

$$\mathcal{L}_{+,-}^A \cong \exp \left(-2t^2 \sum_{k \in \mathbb{G}_A} A_k^2 \cos^2 \frac{E_k t}{2} \text{sinc}^2 \frac{E_k t}{2} \right), \quad (37a)$$

$$\mathcal{L}_{+,-}^B \cong \exp \left(-\frac{1}{2} t^4 \sum_{k \in \mathbb{G}_B} E_k^2 B_k^2 \text{sinc}^4 \frac{E_k t}{2} \right). \quad (37b)$$

As the number of pseudo-spins is large, the details of individual pseudo-spins are not important to the decoherence but what matters is the excitation spectrum, which is defined as⁸

$$S_A(\varepsilon) \equiv \sum_{k \in \mathbb{G}_A} \delta(\varepsilon - E_k) A_k^2, \quad (38a)$$

$$S_B(\varepsilon) \equiv \sum_{k \in \mathbb{G}_B} \delta(\varepsilon - E_k) B_k^2, \quad (38b)$$

for the pair-flips in group \mathbb{G}_A and \mathbb{G}_B , respectively. The decoherence in terms of the excitation spectra is

$$\mathcal{L}_{+,-}^A \cong \exp \left(-2t^2 \int d\varepsilon S_A(\varepsilon) \cos^2 \frac{\varepsilon t}{2} \text{sinc}^2 \frac{\varepsilon t}{2} \right), \quad (39a)$$

$$\mathcal{L}_{+,-}^B \cong \exp \left(-\frac{1}{2} t^4 \int d\varepsilon S_B(\varepsilon) \varepsilon^2 \text{sinc}^4 \frac{\varepsilon t}{2} \right). \quad (39b)$$

We are now ready to analyse some important features of the FID, including the short-time behaviour, the dependence on the field strength and on the QD size, and the emergence of Markovian decay.

In the timescale $t \lesssim E_k^{-1}$, the pair-flips cannot be described by energy-conserving scattering events, but should be understood in terms of quantum evolution. The decoherence is a highly non-Markovian dynamics. Particularly, in the very initial stage ($t \ll E_k^{-1}$), according to equation (36), the electron spin coherence is well approximated as

$$\mathcal{L}_{+,-}^A \approx \exp \left(-t^2 / T_{2,A}^2 \right), \quad (40a)$$

⁸ The definition of the excitation spectrum at vanishing frequency, however, is tricky as the density of states at $\varepsilon = 0$ can be made arbitrarily large by enclosing nuclear spins out of the range of the electron wavefunction, which contribute to local pair-flips costing zero hyperfine energy ($E_k = 0$) and inducing no decoherence. To remove such indefiniteness, we indeed define the regularized excitation spectra to be $\tilde{S}_{A/B}(\varepsilon) \equiv \int_{\varepsilon - \varepsilon_0}^{\varepsilon + \varepsilon_0} \omega^2 S_{A/B}(\omega) d\omega / \int_{\varepsilon - \varepsilon_0}^{\varepsilon + \varepsilon_0} \omega^2 d\omega$, where ε_0 is chosen such that $t^{-1} \gg \varepsilon_0 > B_k$.

$$\mathcal{L}_{+,-}^B \approx \exp(-t^4/T_{2,B}^4), \quad (40b)$$

which is not an exponential decay, a typical indicator of Markovian dynamics. Note that the exponential indices are substantially different from the semiclassical theory predictions [25, 26]. The decoherence times in the short-time limit are

$$T_{2,A} \approx \left(2 \sum_{k \in \mathbb{G}_A} A_k^2\right)^{-1/2} \sim \frac{N\Omega_e}{\mathcal{A}_\alpha^2}, \quad (41a)$$

$$T_{2,B} \approx \left(\frac{1}{2} \sum_{k \in \mathbb{G}_B} E_k^2 B_k^2\right)^{-1/4} \sim \frac{N^{5/12}}{B_k^{1/2} \mathcal{A}_\alpha^{1/2}}. \quad (41b)$$

Here, we have used the facts that $A_k \sim N^{-2} \mathcal{A}_\alpha^2 \Omega_e^{-1}$, the number of spin pairs connected by the extrinsic hyperfine-mediated interaction is about N^2 , the number of spin pairs connected by the intrinsic interaction is of the order of N , and the typical hyperfine energy cost E_k for spin flip–flops between neighbouring nuclei is of the order of $N^{-1/3} a_{n,\alpha} \sim N^{-4/3} \mathcal{A}_\alpha$.

A few features of the FID in the short-time limit ($t \ll E_k^{-1}$) are summarized as follows. The decoherence caused by the intrinsic interaction has a quartic exponential decay profile and that by the hyperfine-mediated interaction has a quadratic one. So the decoherence is initially dominated by the hyperfine-mediated interaction and then cross over to the regime dominated by the intrinsic interaction as time increases. For a QD with 10^5 nuclear spins and an external field of strength 10 tesla, the decoherence times due to the local and nonlocal pair-flips, T_B and T_A , are both of the order of $1 \mu\text{s}$. The two decoherence times have very different dependence on the QD size (measured by N) and the field strength (measured by Ω_e), as can be seen from equation (41). The hyperfine-mediate interaction could be the dominating decoherence mechanism when the QD size is small or when the external magnetic field is weak (e.g. $B_{\text{ext}} \sim 1$ tesla). The intrinsic interaction becomes dominating for large QD or when the field is strong (see figure 7). In a QD of proper size and under an external field of proper strength, the crossover from the quadratic to the quartic exponential decay presents in a visible regime, i.e. in a time range where the electron spin decoherence has decayed by a finite amount but not vanished yet (see the curves for $B_{\text{ext}} = 12$ tesla in figure 7(b)). The crossover can be tuned to occur after the coherence has vanished (as for $B_{\text{ext}} < 10$ tesla in figure 7) or before the decoherence is significant (as for $B_{\text{ext}} > 14$ tesla for the QD in figure 7, in which the data are not shown), then the decoherence in the visible regime will be dominated by the hyperfine-mediated interaction or the intrinsic interaction, respectively. Particularly, in the strong field regime, the decoherence time of FID in single-system dynamics would be greatly underestimated without the intrinsic nuclear spin interaction included [17]–[22].

In general, the Markovian decay will emerge with time increasing, particularly for $t > E_k^{-1}$, when the Fermi-Golden rule starts to come into effect (as indicated by the sinc function in equations (37) and (39)). Actually, the spin coherence at time t is mostly determined by the pseudo-spins with energy cost $E_k < t^{-1}$, while the effects of pseudo-spins with higher precession frequency are cancelled out by destructive interference (between the fast-oscillating sinc functions). In the long-time limit (while $t \ll B_k^{-1}$ is still satisfied), the excitation spectra $S_{A/B}(\varepsilon)$ in equation (39) can be taken as ‘flat’ in the frequency range $[-1/t, 1/t]$, which covers those pseudo-spins that contribute significantly to the decoherence. The assumption of ‘flat’




Figure 7. (a) Electron spin coherence as functions of time for various field strengths. (b) The logarithm plot of (a), in which the curve for $B_{\text{ext}} = 12$ tesla is compared with the contribution by the hyperfine-mediated interaction (the dashed line) and that by the intrinsic interaction (the dotted line), respectively. The size of the InAs dot is $33 \times 33 \times 3 \text{ nm}^3$ and the nuclear-spin initial state $|\mathcal{J}\rangle$ is randomly selected from an ensemble at temperature 1 K. The field strength is indicated by the numbers for each curve.

spectra is just the Markovian approximation. So for time much greater than the inverse widths of the excitation spectra (which are roughly the inverse typical energy cost of the pair-flips, E_k^{-1}), the electron spin coherence is approximated as

$$\mathcal{L}_{+,-}^A \approx e^{-tS_A(t^{-1}) \int 4 \cos^2 x \text{sinc}^2 x dx} \equiv e^{-t/T_{\infty,A}}, \quad (42a)$$

$$\mathcal{L}_{+,-}^B \approx e^{-tS_B(t^{-1}) \int 4 \sin^2 x \text{sinc}^2 x dx} \equiv e^{-t/T_{\infty,B}}, \quad (42b)$$

presenting an exponential decay, a signature of the Markovian dynamics. The decoherence rate in the Markovian regime is determined by the weighted density of states of the elementary excitations, i.e. the pair-flips, which, by the definition of the excitation spectra in equation (38), is related to the Fermi–Golden rule by

$$T_{\infty,A}^{-1} = \xi_A \sum_k A_k^2 \delta(t^{-1} - E_k), \quad (43a)$$

$$T_{\infty,B}^{-1} = \xi_B \sum_k B_k^2 \delta(t^{-1} - E_k), \quad (43b)$$

where $\xi_{A/B}$ denotes the integral in equation (42) and is of the order of unity.




Figure 8. (a) The electron spin coherence in a QD of size $15 \times 15 \times 2.4 \text{ nm}^3$, under a field of 10 tesla and temperature of 1 K. The lower curves are calculated with the hyperfine-mediated pair-flips neglected, and the dotted lines are the short-time profile. The non-Markovian-to-Markovian crossover is observed. (b) The excitation spectra for the nonlocal and local pair-flips.

Under the conditions considered in this paper, the Markovian behaviour will not be fully developed before the coherence has totally vanished, but instead a crossover behaviour, namely, a reduction of the exponential index could be observed in the visible regime for certain QD size and field strength. An example of such non-Markovian-to-Markovian crossover is shown in figure 8(a). The typical energy cost of local pair-flips and that of nonlocal ones is in the order of $\mathcal{A}_\alpha N^{-4/3}$ and $\mathcal{A}_\alpha N^{-1}$, respectively, so the excitation spectrum of local pair-flips is much narrower than that of nonlocal ones (as shown in figure 8(b)). Thus we expect that the Markovian dynamics will emerge earlier for the quadratic exponential decay (caused by nonlocal pair-flips) than for the quartic one (caused by local pair-flips). To observe the non-Markovian-to-Markovian crossover before the decoherence is complete, one should have $E_k^{-1} < T_{2,A/B}$ which is satisfied in relatively small QDs (for decoherence contributed by local pair-flips) or under relatively strong external field (for decoherence contributed by nonlocal pair-flips).

Due to the rich crossover behaviours discussed above, the decoherence in general cannot be characterized by a single time parameter T_2 . We introduce a decoherence time $T_{1/e}$ to quantify the time when the electron spin coherence has decayed to $1/e$ of its initial value. The decoherence time $T_{1/e}$ can approach $T_{2,A}$, $T_{2,B}$, $T_{\infty,A}$, or $T_{\infty,B}$, depending on the QD size and the field strength which set the conditions for the crossovers. The field- and the size-dependence of different decoherence times shown in figures 9 and 10, respectively, are consistent with the analysis above. For comparison, we also plot the decoherence time measured by the Hahn echo T_H , which will be studied in the next section.




Figure 9. Field dependence of decoherence times with the inhomogeneous broadening effect excluded ($T_{1/e}$ —time for FID coherence being $1/e$ of its initial value, $T_{2,A}$ —FID decoherence time resulting solely from hyperfine-mediated pair-flips, $T_{2,B}$ —FID decoherence time resulting solely from the intrinsic nuclear spin interaction, T_H —decay time of the Hahn echo signal). The $\sqrt{2}T_{2,B}$ is plotted to compare with the Hahn echo decay time. The QD is as in figure 7.




Figure 10. QD-size dependence of decoherence times with the inhomogeneous broadening excluded (see figure 9 and the text for definition). The $\sqrt{2}T_{2,B}$ is plotted to compare with the Hahn echo decay time. The QD size is varied with fixed width : depth : height ratio 33 : 33 : 6, the field strength is 10 telsa, and the temperature is 1 K.

4. Hahn echo signal

The FID studied in the previous section is not visible in ensemble experiments in which dephasing due to inhomogeneous broadening is faster by orders of magnitude than decoherence by dynamical entanglement. It might be difficult in the near future to directly observe the

single-system dynamics by filtering out the inhomogeneous broadening with projective measurement of the local Overhauser field [20], [67]–[69]. Thus the spin echo may be used to study the electron spin decoherence problem in QDs. In Hahn echo experiments, the static Overhauser field experienced by the electron spin effectively changes its sign each time the electron spin is flipped by a short π -pulse. Considering the simplest configuration in which only one pulse is applied at $t = \tau$, the precession phase accumulated from the random local fields is eliminated at $t = 2\tau$. Thus the decay of Hahn echo signals at $t = 2\tau$ is solely due to the dynamical entanglement. The decay time of Hahn echo signals is usually used to quantify the spin decoherence time. We shall show, however, the Hahn echo decay time cannot be equated with the decoherence time, as the decoherence by hyperfine-mediated interaction will be virtually eliminated from the echo signal and that by intrinsic interaction is also suppressed to some extent.

The π -rotation or flip of the electron spin for Hahn echo can be operated by a GHz microwave pulse [71]–[73], which is only marginally fast enough for eliminating the rapid dephasing by inhomogeneous broadening in III–V compound QDs. In optical control of an electron spin [74], an arbitrary rotation of the electron spin can be completed in the timescale of 10 ps via exciton-mediated Raman processes. With respect to the timescale of the electron spin decoherence, the optical pulses can be considered instantaneous. The recent experiment on double GaAs QDs also employs a rather long dc voltage pulse to control the singlet–triplet transition to realize the spin echo [40], which does not satisfy the instantaneous pulse condition. It is certainly interesting to study the electron spin decoherence under the control of finite-duration pulses [56], but in this paper we would rather focus on the case of instantaneous pulses.

As our general rule, the Hahn echo signal is determined by the distinguishability δ_k^2 between the conjugate pseudo-spins. In the rotating reference frame rested on the electron spin, the pseudo-spin states after the π -pulse applied at τ are

$$|\psi_k^\pm(t > \tau)\rangle = \hat{U}_k^\mp(t - \tau)\hat{U}_k^\pm(\tau)|\downarrow\rangle, \quad (44)$$

which is equivalent to the conjugate pseudo-spin states $|\psi_k^\pm(t)\rangle$ exchanging their pseudo-fields χ_k^\pm when the electron spin is flipped. The trajectories of the Bloch vectors, projected to the x – y -plane, are shown in figure 11, both for local and for nonlocal pair-flips. The pseudo-spin states after the flip pulse are

$$|\psi_k^\pm(\tau + t')\rangle = \left[\cos \frac{\chi_k^\mp t'}{2} \cos \frac{\chi_k^\pm \tau}{2} - \sin \frac{\chi_k^\mp t'}{2} \cdot \sin \frac{\chi_k^\pm \tau}{2} - i\hat{\sigma}_k \cdot \left(\cos \frac{\chi_k^\mp t'}{2} \sin \frac{\chi_k^\pm \tau}{2} + \cos \frac{\chi_k^\pm \tau}{2} \sin \frac{\chi_k^\mp t'}{2} + \sin \frac{\chi_k^\mp t'}{2} \times \sin \frac{\chi_k^\pm \tau}{2} \right) \right] |\downarrow\rangle. \quad (45)$$

The Hahn echo signal is

$$\mathcal{L}_{+,-}^s(2\tau) = \prod_k |\langle \psi_k^-(2\tau) | \psi_k^+(2\tau) \rangle|. \quad (46)$$

With the same arguments as in section 3, the electron spin decoherence can be factorized into two contributions, namely $\mathcal{L}_{+,-}^s = \mathcal{L}_{+,-}^A \times \mathcal{L}_{+,-}^B$ with

$$\mathcal{L}_{+,-}^{A/B} \approx \prod_{k \in \mathbb{G}_{A/B}} \exp(-\delta_k^2/2). \quad (47)$$




Figure 11. (a) The trajectories (projected to the x - y -plane) of the conjugate pseudo-spins by the intrinsic interaction. The conjugate pseudo-spins $|\psi_k^\pm\rangle$ exchange their pseudo-fields χ_k^\pm at $t = \tau$ when the electron spin is flipped by a short pulse. The solid and dotted curves denote the trajectories before and after the flipping pulse, respectively. (b) The same as (a) but for a nonlocal pair-flip by the hyperfine-mediated interaction.

For the local pair-flips driven by the intrinsic nuclear spin interaction, the distinguishability between conjugate pseudo-spins at $t = 2\tau$ is calculated by setting $D_k = A_k = 0$:

$$\delta_k^2 \cong \frac{1}{4}(2\tau)^4 E_k^2 B_k^2 \text{sinc}^4 \frac{E_k \tau}{2} \approx \frac{1}{4}(2\tau)^4 E_k^2 B_k^2, \quad (48)$$

where the second approximation holds for $\tau \ll E_k^{-1}$. The short-time behaviour is

$$\mathcal{L}_{+,-}^B(2\tau) \approx e^{-(2\tau)^4/T_{H,B}^4}, \quad (49)$$

the same as the FID signal except that the decay time

$$T_{H,B} = \sqrt{2}T_{2,B}. \quad (50)$$

For nonlocal pair-flips driven by the hyperfine-mediated interaction, the pseudo-fields χ_k^\pm are opposite to each other if the energy cost due to the diagonal nuclear spin interaction is neglected ($D_k = 0$). In this case, the pseudo-spins simply reverse their precession directions when the electron spin is flipped, returning to its original states at the echo time, as shown in figure 11(b), so the decoherence by the hyperfine-mediated pair-flips is fully eliminated if D_k is negligible. The elimination of the hyperfine interaction induced decoherence by spin echo was noticed before in numerical simulations of few-spin systems [23]. As the leading order of the distance δ_k vanishes at the echo time, the secondary effect of the diagonal nuclear spin interaction becomes important. The distinguishability between conjugate pseudo-spins for nonlocal pair-flips, including the effect of D_k , is given by

$$\delta_k^2 \cong \frac{1}{4}(2\tau)^4 D_k^2 A_k^2 \text{sinc}^4 \frac{E_k \tau}{2} \approx \frac{1}{4}(2\tau)^4 D_k^2 A_k^2, \quad (51)$$

where again, the second approximation holds for $\tau \ll E_k^{-1}$. The residual decoherence due to the interplay of the hyperfine-mediated interaction and the diagonal nuclear spin interaction has the quartic exponential form for $\tau \ll E_k^{-1}$ as

$$\mathcal{L}_{+,-}^A(2\tau) \cong e^{-(2\tau)^4/T_{H,A}^4}. \quad (52)$$

The corresponding decoherence time is

$$T_{H,A} = \left(\sum_{k \in \mathbb{G}_A} D_k^2 A_k^2 / 8 \right)^{-1/4} \sim \frac{N^{1/2} \Omega_e^{1/2}}{\mathcal{A}_\alpha D_k^{1/2}}, \quad (53)$$

proportional to the square root of the field strength.

Now we compare the echo signals to the FID signal in single-system dynamics (see figures 9 and 12). The observation is that the decoherence due to the dynamical quantum entanglement in general is not measured by the Hahn echo signals. Firstly, the FID and the echo signals could have qualitatively different short-time behaviours, as the former could have quadratic exponential decay due to the hyperfine-mediated interaction while the latter always has the quartic exponential decay. Secondly, the decay timescale could be dramatically different from the FID to the Hahn echo. The difference in the decay time can be discussed in three regimes of field strength, roughly divided by

$$\text{I} : \Omega_e \lesssim \mathcal{A}_\alpha N^{-1/6}, \quad (54a)$$

$$\text{II} : \mathcal{A}_\alpha N^{-1/6} \ll \Omega_e \lesssim \mathcal{A}_\alpha^{3/2} N^{-7/12} B_k^{-1/2}, \quad (54b)$$

$$\text{III} : \mathcal{A}_\alpha^{3/2} N^{-7/12} B_k^{-1/2} \ll \Omega_e, \quad (54c)$$

corresponding to

$$\text{I} : T_{H,B} \gtrsim T_{H,A}, \quad (55a)$$

$$\text{II} : T_{H,A} \gg T_{H,B} \sim T_{2,B} \gtrsim T_{2,A}, \quad (55b)$$

$$\text{III} : T_{2,A} \gg T_{2,B}. \quad (55c)$$

In regime I, the hyperfine-mediated interaction is important both in the FID and in the Hahn echo decay. As the excitation of nonlocal pair-flips is eliminated in the leading order at the echo time (see figure 11(b)), the echo decay time is much longer than the FID decoherence time, namely,

$$T_{2,A} / T_{H,A} \sim \sqrt{N \Omega_e D_k / \mathcal{A}_\alpha^2} \ll 1, \quad (56)$$

for QDs of reasonable sizes. The pair-correlation approximation, however, is not valid in regime I, since the condition that the number of pair-flips in the timescale of $T_{H,A}$ be much fewer than the number of nuclei requires $N \gg \mathcal{A}_\alpha^{4/3} \Omega_e^{-2/3} D_k^{-2/3}$ for the hyperfine mediated nonlocal pair-flips and $N \ll \mathcal{A}_\alpha^{4/3} \Omega_e^{-2/3} D_k^{-2/3}$ for the local pair-flips driven by the intrinsic interaction (see appendix C for estimation of the pair-flip numbers), which cannot be simultaneously satisfied. Theories beyond the pair-correlation approximation need to be developed to explore this regime. In regime II defined by equation (54b), the hyperfine-mediated interaction contributes significantly to the FID but virtually nothing to the Hahn echo decay. In this regime, the decay time measured in echo signals is much longer than the FID decoherence time as

$$T_{H,B} = \sqrt{2} T_{2,B} \gg T_{2,A}. \quad (57)$$




Figure 12. (a) Comparison of the Hahn echo (dashed green line) and the FID (solid red line) signals. The FID signal is also shown with the hyperfine-mediated pair-flips neglected (dotted blue line). (b) The logarithm plot of (a). The QD is as in figure 7, with $B_{\text{ext}} = 2$ tesla.

In regime III, the hyperfine-mediated interaction is strongly suppressed, so the FID and the Hahn echo decay, both determined by the intrinsic nuclear spin interaction, have essentially the same decay profile but different decay times (differing by a factor of $\sqrt{2}$).

As the leading order effect of the hyperfine-mediated interaction is eliminated by spin echo, it is critical to include the intrinsic nuclear spin interaction in studying the spin echo signals. Note even for relatively weak external field ($B_{\text{ext}} < 0.01$ T in regime I for typical QDs) where the decoherence induced by the intrinsic interaction is relatively slow ($T_{H,B} > T_{H,A}$), the intrinsic interaction is still important to give the correct residual decoherence effect together with the hyperfine-mediated interaction (see equation (51)). Actually, it has been shown before that if the intrinsic nuclear spin interaction is neglected and the T_1 (electron spin-flip) process is suppressed, the decoherence caused by the hyperfine interaction is totally removed by spin echo [23] (which can be easily understood in the second-order term used in this paper by noticing that $\hat{H}_+ = -\hat{H}_-$ in equation (3) if the intrinsic interaction \hat{H}_B and \hat{H}_D are set zero). Thus it may be concluded that only when the external field is so weak that the T_1 relaxation is significant ($B_{\text{ext}} < T_2^{*-1} \sim 0.001$ T for typical QDs [4]), can the theories neglecting the intrinsic nuclear spin interaction [17]–[22] be applied to study spin echo signals. It should also be mentioned that even in the spin echo signals, the hyperfine-mediated interaction plays a nontrivial role in the relatively weak field regime and our results for spin echo signals reduce to those calculated without the hyperfine-mediated interaction [7, 8] only in regimes II and III.

We have seen notable difference between the FID and the Hahn echo decay in all the three field strength regimes discussed above. Such difference is indeed a consequence of the modified

dynamics of a mesoscopic bath (the nuclear spins) under the manipulation of the quantum object (the electron spin). In semiclassical spectral diffusion theories, the electron spin experiences passively the “background” of fluctuating local fields. In the full quantum theory, the electron spin dynamics actively alters the mesoscopic bath dynamics (as the nuclear spin pair-flips depends dramatically on the electron spin state). Such a view paves the path toward manipulation of the nuclear spin dynamics in mesoscopic QDs and control of the electron spin decoherence.

The active modification of the bath dynamics by the manipulation of the quantum object also manifests itself in the effect of the diagonal nuclear spin interaction on the Hahn echo signal (see equation (51)). The diagonal nuclear spin interaction contributes much less energy cost of a pair-flip than the hyperfine interaction ($D_k \ll E_k$) and has no direct effect on the effective Overhauser field felt by the electron spin. Thus it is expected that the diagonal interaction terms should have little effect on the electron spin decoherence. The calculation of the FID signal in section 3 indeed confirms such a semiclassical argument. The difference between the quantum theory and its semiclassical counterparts becomes significant when the electron spin is under control. In semiclassical theories, the electron spin only passively detects the fluctuating local field which is essentially independent of the electron dynamics, so the diagonal nuclear spin interaction should contribute no more than it does in the FID configuration. In the quantum theory, the nuclear spin dynamics conditioned on the electron spin state is held responsible for the electron spin decoherence. The diagonal nuclear spin interaction affects the nuclear spin dynamics significantly and thus contributes to the decoherence when the nuclear spin dynamics is under active control by the manipulation of the electron spin state. In the next sections, we will see the importance of the diagonal nuclear spin interaction in determining some qualitative decoherence features.

5. Disentanglement and recoherence

In the previous section, we have seen that the decoherence caused by the hyperfine-mediated interaction is nearly eliminated in Hahn echo signals. Such elimination of decoherence is realized because the hyperfine-mediated pair-flips are reversed when the electron spin is flipped at $t = \tau$ (see figure 11(b)) and hence the pseudo-spin states $|\psi_k^\pm(t)\rangle$ return to their original positions at $t = 2\tau$. As the decoherence is caused by the entanglement between the electron spin and the nuclear spins, the elimination of decoherence, also called ‘recoherence’, can be understood in terms of disentanglement. For nonlocal pair-flips under Hahn echo control, the disentanglement is realized at $t = 2\tau$ when the electron spin and the pseudo-spin are brought back to a factorized state $C_+|+\rangle|\psi_k^+(2\tau)\rangle + C_-|-\rangle|\psi_k^-(2\tau)\rangle$ with $|\psi_k^\pm(2\tau)\rangle = |\psi_k^\pm(0)\rangle = |\downarrow\rangle$.

In general, a pseudo-spin (or the environment) is disentangled from the electron spin (or the quantum system in contact with the environment) whenever its states $|\psi_k^\pm(t)\rangle$ for different electron spin states $|\pm\rangle$ become identical, but need not return to the original states $|\psi_k^\pm(0)\rangle$. When the electron spin is flipped, the conjugate pseudo-spins exchange their pseudo-fields, so their trajectories on the Bloch sphere will inevitably intersect, leading to the disentanglement⁹. The intersection of conjugate pseudo-spins driven by the intrinsic nuclear spin interaction is

⁹ A pseudo-spin is disentangled from the electron spin after a full precession cycle in free evolution. But the pseudo-field strength χ_k^\pm is different for different pseudo-spins, so is the disentanglement time, making it impossible for the recoherence to occur in QDs of many nuclear spins without applying a controlling pulse. For small QDs, however, spontaneous recovery of coherence in FID may be observed.

actually seen in figure 11(a), which, unlike that for nonlocal pair-flips, occurs at a time different from the echo time. The intersection time, in general, is different for different pair-flips, making it impossible to fully recover the lost electron spin coherence. Nonetheless, in the short-time limit ($\tau \ll E_k^{-1}$), the distance between conjugate pseudo-spins is eliminated in the leading order of time, simultaneously for the same group of pair-flips, as have been seen for nonlocal pair-flips. It is straightforward to verify that for the local pair-flips, such coincident disentanglement does take place at $t = \sqrt{2}\tau$ in the single-pulse Hahn echo configuration (the magic number $\sqrt{2}$ can be understood by noticing that $\delta_k \propto \tau^2$ at short-time limit and $(\sqrt{2}\tau)^2 - \tau^2 = \tau^2$).

With the leading order contribution vanishing, the distinguishability between conjugate pseudo-spins at the disentanglement time ($\sqrt{2}\tau$), including the effect of the diagonal nuclear spin interaction, in the next leading order of τ for local pair-flips is

$$\delta_k(\sqrt{2}\tau) \sim E_k B_k D_k \tau^3. \quad (58)$$

Note that the residual decoherence would otherwise be of a higher order in τ if D_k were set to zero:

$$\delta_k(\sqrt{2}\tau) \sim E_k B_k^3 \tau^4. \quad (59)$$

Here we see again that the small effect of the diagonal nuclear spin interaction (the D_k term) emerges when the nuclear spin dynamics is under control.

A semiclassical spectral diffusion theory also predicts a coherence recovery at a time earlier than the echo time (2τ) when the random field memory time (τ_c) is comparable to the pulse delay time [75]. But the semiclassical theory differs qualitatively from the quantum counterpart on at least two aspects: (i) the recovery time depends on τ_c , being about $\tau + \tau_c$ for $\tau_c < \tau$ and approaching 2τ when $\tau_c \gg \tau$; and (ii) the exponential index of the decoherence profile in the spectral diffusion theory is essentially unchanged by the control. The latter difference points directly to a fundamental shortness of the semiclassical picture: it does not take into account the active control of the local field fluctuation by the flip of the quantum object but considers the quantum evolution of the bath as a fluctuating field experienced passively by the quantum object.

Inclusion of contributions from both local and nonlocal pair-flips fields the pulse-controlled decoherence in the short-time limit,

$$\mathcal{L}_{+,-}^s(\sqrt{2}\tau) \cong e^{-(\sqrt{2}\tau)^6/T_{\text{eff},B}^6} \times e^{-(\sqrt{2}-2)^2\tau^2/T_{2,A}^2}, \quad (60)$$

where the effective decoherence time due to the residual entanglement from local pair-flips is defined by

$$T_{\text{eff},B}^{-6} \sim \sum_{k \in \mathbb{G}_B} E_k^2 B_k^2 D_k^2. \quad (61)$$

The electron coherence is recovered at $\sqrt{2}\tau$, even when the pulse is applied after the coherence has been totally lost in FID (for $\tau > T_{2,B}$). Such recoherence in single-system dynamics and the spin echo in ensemble dynamics have different physical bases: the former is due to quantum disentanglement, but the latter is due to classical refocusing of random phases. The difference between the two is already evidenced by their different occurrence time. Figure 13(a) plots an example of the real-time dependence of the electron spin coherence under a single-pulse control, which demonstrates the recovery of the coherence even when the FID signal has vanished for the chosen delay time. The contribution from the hyperfine-mediated pair-flips, which are not reversed at $\sqrt{2}\tau$, makes the coherence to be only partially recovered. Under a stronger




Figure 13. (a) The electron spin coherence under the control of a short π -pulse applied at $\tau = 2 \mu\text{s}$ (when the FID signal has vanished), the recoherence at $\sqrt{2}\tau$ is pronounced while no signal survives at the echo-time 2τ . (b) The electron spin coherence under the control of a sequence of evenly spaced pulses. (c) The electron spin coherence under the control of a Carr–Purcell pulse sequence. The arrows indicate positions of the short pulses. The solid blue (dotted red) lines are calculated with(out) including the hyperfine-mediated pair-flips. The QD is as in figure 7 with $B_{\text{ext}} = 10$ tesla. No ensemble average is calculated.

magnetic field, a full recovery at $\sqrt{2}\tau$ can be realized as the hyperfine-mediated interaction is fully suppressed (not shown).

As illustrated in figure 14(a), a sequence of pulses can force the trajectories of conjugate pseudo-spins to cross into each other again and again. In the short-time limit (the delay time $\tau \ll E_k^{-1}$), an equally spaced pulse sequence eliminates the entanglement due to the local pair-flips up to the τ^2 terms, leading to a sequence of recoherence at

$$t_n = \sqrt{n(n+1)}\tau, \quad n = 1, 2, \dots \quad (62)$$

One can verify these magic numbers by using the quadratic dependence of δ_k on time and checking that $(\sqrt{n(n+1)}\tau)^2 - (n\tau)^2 = (n\tau)^2 - (\sqrt{(n-1)n}\tau)^2$. The decoherence caused by the hyperfine-mediated pair-flips under the control of evenly spaced pulses is bounded, which can be understood from the reversed precession of the corresponding pseudo-spins. Figure 13(b) confirms our above analysis. By using a long sequence of pulses at short intervals, the electron spin coherence can be preserved for an arbitrarily long time until other mechanisms of decoherence (such as phonon scattering) come into effect.

The recoherence in the control configurations discussed above, however, is not observable in ensemble experiments, or in theories on ensemble dynamics [7, 8], since the rapid dephasing




Figure 14. Trajectories of conjugate pseudo-spins from the intrinsic nuclear spin interaction (a) under the control of a sequence of pulses equally spaced and (b) under the control of a Carr–Purcell two-pulse sequence. The green bars on the time axis represent the short pulses flipping the electron spin.

due to inhomogeneous broadening will prevent any coherence being observed except around the spin-echo time. It is desirable to design a pulse sequence to force the recoherence from disentanglement to coincide with a spin-echo from phase-refocusing so that the disentanglement effect can be studied in ensemble experiments. This possibility can be seen from figure 14(b). When the electron spin is flipped by two short pulses, for example, the disentanglement time after the second pulse can be adjusted by changing the delay time between the two pulses. It is shown straightforwardly that when the pulse sequence is designed such that the first pulse is applied at $t = \tau$ and the second at $t = 3\tau$, the disentanglement coincides with the echo time at $t = 4\tau$ (by virtue of the change in δ_k , $(4\tau)^2 - (3\tau)^2 - [(3\tau)^2 - \tau^2] + \tau^2 = 0$). Obviously, the hyperfine-mediated nonlocal pair-flips are disentangled from the electron spin at the echo time. A two-pulse sequence so designed happens to be the famous Carr–Purcell sequence, widely used in NMR experiments to dynamically decouple the nuclear spins. The disentanglement studied here, however, is fundamentally different from the dynamical decoupling, as will be discussed later in the next section.

In the rotating frame of the electron spin, the pseudo-spin states at the echo time after a Carr–Purcell sequence is

$$|\psi_k^\pm(4\tau)\rangle = e^{-i\theta_0^\pm \cdot \hat{\sigma}/2} e^{-i\theta_0^\mp \cdot \hat{\sigma}} e^{-i\theta_0^\pm \cdot \hat{\sigma}/2} |\downarrow\rangle = e^{-i\theta_1^\mp \cdot \hat{\sigma}/2} e^{-i\theta_1^\pm \cdot \hat{\sigma}/2} |\downarrow\rangle = e^{-i\theta_2^\pm \cdot \hat{\sigma}/2} |\downarrow\rangle, \quad (63)$$

with the series of angles of rotation defined by

$$\theta_0^\pm \equiv \chi_k^\pm \tau, \quad (64a)$$

$$e^{-i\theta_{l+1}^\pm \cdot \hat{\sigma}/2} \equiv e^{-i\theta_l^\mp \cdot \hat{\sigma}/2} e^{-i\theta_l^\pm \cdot \hat{\sigma}/2}, \quad (64b)$$

The vector θ_l^\pm has a geometrical interpretation as the pseudo-spin effective precession angle θ_l^\pm about the axis along θ_l^\pm at τ , 2τ , and 4τ for $l = 0$, 1 and 2, respectively (the pseudo-spin index k is understood where no confusion is caused). Equation (64b) yields a geometrical recursion

relation:

$$\sin \frac{\theta_{l+1}^{\pm}}{2} = \cos \frac{\theta_l^{\mp}}{2} \sin \frac{\theta_l^{\pm}}{2} + \cos \frac{\theta_l^{\pm}}{2} \sin \frac{\theta_l^{\mp}}{2} + \sin \frac{\theta_l^{\mp}}{2} \times \sin \frac{\theta_l^{\pm}}{2}, \quad (65a)$$

$$\cos \frac{\theta_{l+1}^{\pm}}{2} = \cos \frac{\theta_l^{\mp}}{2} \cos \frac{\theta_l^{\pm}}{2} - \sin \frac{\theta_l^{\mp}}{2} \sin \frac{\theta_l^{\pm}}{2}. \quad (65b)$$

The distinguishability between conjugate pseudo-spins at the echo time 4τ in the leading order of delay time is

$$\delta_k^2 \cong 2^4 \tau^6 D_k^2 (B_k E_k - A_k D_k)^2, \quad (66)$$

for $\tau \ll E_k^{-1}$. This leads to an $\exp(-\alpha\tau^6)$ decay profile which has also been independently discovered by the Das Sarma group [8]. Note the decoherence would otherwise be of a higher order in τ if D_k is set to zero,

$$\delta_k^2 \cong 2^{10} \tau^8 B_k^6 E_k^2. \quad (67)$$

This demonstrates again the role of the diagonal nuclear spin interaction. We note that the decoherence profile in the two-pulse control configuration, in particular the exponential index, is qualitatively different from the predictions of semiclassical theories [44] in which the bath dynamics is accounted for by a randomly fluctuating force (without being actively changed by the manipulation of the quantum object) and the resultant decoherence exponential index is basically unchanged by the pulse control. The recoherence at the echo time 4τ in the Carr–Purcell control configuration is clearly demonstrated in figure 13(c), even though the signal at the Hahn echo time 2τ after the first pulse is absent.

6. Concatenated disentanglement

6.1. Formalism and geometrical picture

We have used the iteration of effective precession angles to derive the pseudo-spin states under the Carr–Purcell control in equations (63)–(65). Such an iteration formalism inspires us to borrow the idea of concatenation control in dynamical decoupling recently studied for preserving coherence in quantum computation [54]–[57]. To illustrate the basic idea, we can imagine that if χ_k^{\pm} in equation (64a) is replaced by the effective field corresponding to the precession angles at the Carr–Purcell echo time, i.e., θ_2^{\pm}/τ , the distance between the conjugate pseudo-spins would be eliminated to an even higher order of the delay time. Such a replacement can be done recursively, and a concatenated pulse sequence can be so designed to control the decoherence. The unit cell of pulses to be concatenated does not have to be the Carr–Purcell sequence, but it could be either simpler or more complicated, or even various types of unit sequences can be interwoven to construct a sophisticated concatenation.

To exemplify the concatenated disentanglement for decoherence control, we choose the simplest concatenation sequence which is recursively defined by equation (64). Figure 15 illustrates how such a sequence can be concatenated to any order. The 0th order is a FID evolution with no pulse control and the $(l+1)$ th order sequence is constructed by two subsequent l th




Figure 15. Concatenated sequences of short-pulses flipping the electron spin, represented by vertical bars.

sequences with or without one extra flipping pulse inserted in between depending on whether l is even or odd. Accordingly, the evolution propagator is iterated as

$$\hat{U}_0^\pm = e^{-i\chi_k^\pm \cdot \hat{\sigma}_k \tau / 2} \equiv e^{-i\theta_0^\pm \cdot \hat{\sigma} / 2}, \quad (68a)$$

$$\hat{U}_{l+1}^\pm = \hat{U}_l^\mp \hat{U}_l^\pm \equiv e^{-i\theta_{l+1}^\pm \cdot \hat{\sigma} / 2}. \quad (68b)$$

For a pictorial understanding of the concatenated control of decoherence, we rewrite the effective precession angles as

$$\sin \frac{\theta_l^\pm}{2} \equiv \mathbf{n}_l^\pm \equiv \mathbf{R}_l \pm \mathbf{r}_l. \quad (69)$$

Without confusion, the vectors \mathbf{n}_l^\pm will also be referred to as effective precession angles since $\mathbf{n}_l^\pm \approx \theta_l^\pm / 2$ for small precession angles. As depicted in figure 16(a), the conjugate precession angles are decomposed into the common part \mathbf{R}_l and the difference part \mathbf{r}_l . Then, the recursion in equation (65) is rewritten as

$$\mathbf{r}_0 = \frac{1}{2} \sin \frac{\chi_k^+ \tau}{2} - \frac{1}{2} \sin \frac{\chi_k^- \tau}{2}, \quad (70a)$$

$$\mathbf{R}_0 = \frac{1}{2} \sin \frac{\chi_k^+ \tau}{2} + \frac{1}{2} \sin \frac{\chi_k^- \tau}{2}, \quad (70b)$$

$$\mathbf{R}_1 = \cos \frac{\chi_k^- \tau}{2} \sin \frac{\chi_k^+ \tau}{2} + \cos \frac{\chi_k^+ \tau}{2} \sin \frac{\chi_k^- \tau}{2}, \quad (70c)$$

$$\mathbf{r}_l = 2\mathbf{R}_{l-1} \times \mathbf{r}_{l-1}, \quad (l \geq 1), \quad (70d)$$

$$\mathbf{R}_l = 2\mathbf{R}_{l-1} \sqrt{1 - R_{l-1}^2 - r_{l-1}^2}, \quad (l \geq 2). \quad (70e)$$




Figure 16. (a) The effective precession vectors for conjugate pseudo-spins are decomposed into the common component and the difference part. (b) The iteration of the effective precession vectors from the l th order to the $(l+1)$ th order concatenation. (c) The pseudo-fields are decomposed into the perpendicular and the parallel components.

The distance δ_k is in the same order of the difference part. Particularly, for the l th order concatenation, the distance at $\tau_l \equiv 2^l \tau$ is

$$\begin{aligned} \delta_l^2 &= 1 - |\langle \psi_k^- | \psi_k^+ \rangle|^2 = 1 - \left| \left\langle \downarrow \left| \left(\sqrt{1-n_l^2} + \mathbf{n}_l^- \cdot \hat{\sigma} \right) \left(\sqrt{1-n_l^2} - \mathbf{n}_l^+ \cdot \hat{\sigma} \right) \right| \downarrow \right\rangle \right|^2 \\ &= 1 - \left| \left\langle \downarrow \left| 1 - n_l^2 + \mathbf{n}_l^- \cdot \mathbf{n}_l^+ - i2\mathbf{r}_l \cdot \hat{\sigma} \sqrt{1-n_l^2} - i(\mathbf{n}_l^+ \times \mathbf{n}_l^-) \cdot \hat{\sigma} \right| \downarrow \right\rangle \right|^2 \\ &= 4r_l^2 \left[1 - r_l^2 - \left(z_{l+1}R_l - z_l\sqrt{1-R_l^2-r_l^2} \right)^2 \right], \end{aligned} \quad (71)$$

for $l \geq 1$, where $n_l \equiv n_l^+ = n_l^- = \sqrt{R_l^2 + r_l^2}$ and z_l is the z -component of \mathbf{r}_l/r_l .

6.2. Short-time behaviours

To gain some insight into the concatenated control of decoherence, first consider the short-time limit $\tau \ll E_k^{-1}$. There, the recursion of the rotation angles has an intuitive form as

$$\mathbf{R}_l \approx 2^l \mathbf{R}_0 \approx 2^l \tau \frac{\boldsymbol{\chi}_k^+ + \boldsymbol{\chi}_k^-}{4} = (2B_k, 0, D_k) \frac{\tau_l}{2}, \quad (72a)$$

$$\mathbf{r}_0 \approx (\boldsymbol{\chi}_k^+ - \boldsymbol{\chi}_k^-) \frac{\tau}{4} = (2A_k, 0, E_k) \frac{\tau}{2}, \quad (72b)$$

$$\mathbf{r}_l \approx 2^l \mathbf{R}_0 \times \mathbf{r}_{l-1}, \quad (l \geq 1), \quad (72c)$$

which holds until the precession angle R_l approaches the order of unity at a threshold order of concatenation l_0 given by

$$l_0 \approx -\log_2(B_k \tau). \quad (73)$$

The distinguishability between conjugate pseudo-spins in the leading order of delay time is

$$\delta_l^2 \cong 4r_l^2 (1 - z_l^2), \quad (74)$$

determined by the difference between the conjugate precession vectors. The concatenated control can be understood in the geometrical picture shown in figure 16(b). In the FID (the 0th order concatenation), the common part of the precession angles R_0 is roughly determined by the nuclear spin interaction strength times the time, i.e. $B_k \tau$, which is much less than 1 in the timescale of interest, and the difference part r_0 is roughly the precession angle given by the hyperfine energy cost E_k . In the 1st order concatenation, the common part \mathbf{R}_1 is approximately in the same direction as \mathbf{R}_0 with the angle roughly doubled if $E_k \tau \ll 1$, and the difference part \mathbf{r}_1 is perpendicular to both \mathbf{R}_0 and \mathbf{r}_0 with the amplitude reduced by a factor of $2R_0$ from r_0 . Starting from the 2nd order concatenation, the common part \mathbf{R}_l will be along the same direction as the 1st order one \mathbf{R}_1 and the difference part \mathbf{r}_l will be alternatively in the two orthogonal directions \mathbf{r}_1 and \mathbf{r}_2 , both perpendicular to the common part direction \mathbf{R}_l . By each further level of concatenation, the common part is increased by a factor of two and the difference is reduced by a factor of $2R_l$ for $l < l_0$. Thus each level of concatenation will reduce the difference between effective conjugate precession angles by the order of the delay time times the nuclear spin interaction strength and remove the decoherence accordingly, until the controlling effect is saturated at the threshold level l_0 .

With the short-time approximation of the precession angles given in equation (72), the distinguishability between conjugate pseudo-spins in the leading order of τ is

$$\delta_l^2 \cong 4\tau^{2l+2} 2^{l(l-1)} (1 - z_l^2) (4B_k^2 + D_k^2)^{l-1} (B_k E_k - A_k D_k)^2, \quad (75)$$

with

$$z_l^2 = \begin{cases} 0 & (l = 1, 3, \dots), \\ 4B_k^2 (4B_k^2 + D_k^2)^{-1} & (l = 2, 4, \dots). \end{cases} \quad (76)$$

Alternatively, for a fixed time t , a certain order of concatenation control with $\tau = t/2^l$ gives

$$\delta_l^2 \cong 4t^{2l+2} 2^{-l(l+3)} (1 - z_l^2) (4B_k^2 + D_k^2)^{l-1} (B_k E_k - A_k D_k)^2, \quad (77)$$

which can be suppressed to an arbitrary power of the echo time t times the nuclear spin interaction strength under the condition

$$R_l \sim B_k t \ll 1. \quad (78)$$

The electron spin coherence under control is

$$\mathcal{L}_{+,-}^s(\tau_l) \cong \prod_k e^{-\delta_l^2/2} \cong e^{-\tau_l^{2l+2}/T_0^{2l+2}}, \quad (79)$$

decaying exponentially with powers of τ at 2, 4, 6, 8, ..., for the concatenation order 0, 1, 2, 3, ..., respectively.

To show the effect of the diagonal nuclear spin interaction in the decoherence control, we note that in equation (76) $z_{2m} = 1$ if $D_k = 0$ and thus δ_k^2 vanishes in the given order. In this case, the distinguishability between the conjugate pseudo-spins in the next leading order for $l = 2m$ is

$$\delta_l^2 \cong 4R_l^2 r_l^2 \cong \tau^{2l+4} 2^{l(l+3)} E_k^2 B_k^{2l+2}. \quad (80)$$

So when D_k is put to zero, the coherence decays exponentially with powers of τ at 2, 4, 8, 8, 12, 12, ... for concatenation order 0, 1, 2, 3, 4, 5, ..., respectively.

The effective preservation of the coherence by concatenated control is demonstrated by the numerical simulations presented in figure 17. Note that the coherence shown in the figure is at the time $2^l \tau$ which doubles at every order of concatenation. The role of the diagonal part of the nuclear spin interaction is demonstrated (compare figures 17(b) and (c)).




Figure 17. (a) The electron spin coherence under the l th order concatenation control, $l = 0, \dots, 5$, as functions of the pulse delay time τ . (b) The logarithm plot of (a). (c) The same as (b), but with the diagonal nuclear spin interaction put to zero. The QD is as in figure 13.

6.3. Controlling small parameters

The suppression of the decoherence by concatenated control, however, does not depend on the short-time condition $\tau \ll E_k^{-1}$. This point can be seen from equation (70): an iteration of concatenation suppresses the decoherence further as long as the common component of the conjugate precession angles $R_l \ll 1$, without requiring $E_k \tau \ll 1$. Since R_0 is of the order of the nuclear spin interaction strength times the delay time ($B_k \tau$), the condition for a concatenated sequence to be efficient in eliminating the decoherence is that the delay time is much shorter than the inverse strength of the nuclear spin interaction, i.e.

$$\tau \ll B_k^{-1}, D_k^{-1}, A_k^{-1}, \quad (81)$$

which is a much less stringent condition than $t \ll E_k^{-1}$. To calculate the electron spin coherence under this relaxed ‘short-time’ condition, we decompose the conjugate pseudo-fields in such a way that

$$\chi_k^+ = \mathbf{a} + \mathbf{b}, \quad (82a)$$

$$\chi_k^- = \mathbf{a} - \mathbf{c}, \quad (82b)$$

with $\mathbf{a} \perp \mathbf{b} \parallel \mathbf{c}$, as depicted in figure 16(c). Then, we are able to separate different timescales according to the orders of magnitude of different components. We notice that a and $|b - c|$ are of the order of the nuclear spin interaction strength and b and c are of the order of the hyperfine energy cost E_k . Within timescales given by equation (81), the precession angles in the leading order of the nuclear spin interaction strength are

$$\mathbf{r}_0 \cong \sin \frac{(\mathbf{b} + \mathbf{c}) \tau}{4}, \quad (83a)$$

$$\mathbf{R}_0 \cong \frac{(\mathbf{b} - \mathbf{c}) \tau}{4} \cos \frac{(b + c) \tau}{4} + \frac{\mathbf{a} \tau}{2} \operatorname{sinc} \frac{(b + c) \tau}{4}, \quad (83b)$$

$$\mathbf{R}_l \cong 2^{l-2} (\mathbf{b} - \mathbf{c}) \tau + 2^{l-1} \mathbf{a} \tau \operatorname{sinc} \frac{(b + c) \tau}{2}, \quad (l > 0), \quad (83c)$$

$$r_l \cong 2^{l(l-1)/2} \tau^{l+1} \frac{a(b+c)}{4} \operatorname{sinc}^2 \frac{(b+c)\tau}{4} \left[\frac{(b-c)^2}{4} + a^2 \operatorname{sinc}^2 \frac{(b+c)\tau}{2} \right]^{(l-1)/2} \quad (83d)$$

$$\begin{aligned} &\leq 2^{l(l-1)/2} \tau^{l+1} \frac{a}{4} |\chi_k^+ - \chi_k^-| \left| \frac{\chi_k^+ + \chi_k^-}{2} \right|^{l-1} \\ &\cong 2^{\frac{l(l-1)}{2}} \tau^{l+1} |E_k B_k - A_k D_k| (4B_k^2 + D_k^2)^{l-1/2}, \quad (l > 0), \end{aligned} \quad (83e)$$

under the condition $R_l \ll 1$. The z -component of the direction of \mathbf{r}_l is

$$z_l^2 \cong \begin{cases} 0 & (l = 2m + 1), \\ \frac{4B_k^2 \operatorname{sinc}^2(E_k \tau)}{4B_k^2 \operatorname{sinc}^2(E_k \tau) + D_k^2} & (l = 2m + 2). \end{cases} \quad (84)$$

The distinguishability between conjugate pseudo-spins δ_k^2 is determined by equation (71). The decoherence is estimated accordingly.

It is obvious that the small parameter controlling the maximum delay time for a concatenation control to be efficient is the nuclear spin interaction strength $\sqrt{4B_k^2 + D_k^2}$ instead of the object–bath interaction strength (i.e. the hyperfine energy cost E_k). The threshold concatenation level l_0 , beyond which an additional level of concatenation cannot suppress the decoherence further, is determined by the condition that the precession angle approaches 1, i.e. $R_{l_0} \lesssim 1$. With the precession angle given by

$$R_l \cong 2^{l-1} \tau \sqrt{4B_k^2 \operatorname{sinc}^2(E_k \tau) + D_k^2} \leq 2^{l-1} \tau \sqrt{4B_k^2 + D_k^2}, \quad (85)$$

the threshold concatenation level is estimated as

$$l_0 \sim -\log_2 \left(\tau \sqrt{4B_k^2 + D_k^2} \right), \quad (86)$$

also determined by the nuclear spin interaction strength. A close examination of figure 17(a) reveals that the decoherence suppression is still effective even when the pulse delay time τ is substantially longer than the typical value of the inverse hyperfine energy cost E_k^{-1} , which is about $0.5 \mu\text{s}$ for the QD considered.

6.4. Coherence stabilization

The concatenated pulse sequence can be repeated. The propagator of a pseudo-spin for n repeated l th-order concatenation sequences is

$$\hat{U}_{l,n}^{\pm} = \left(\hat{U}_l^{\pm} \right)^n = \exp \left(-in\theta_l^{\pm} \cdot \hat{\sigma}/2 \right). \quad (87)$$

Note that in implementing the pulse sequence for repeated concatenation, a π -pulse should be inserted between two unit sequences if l is an odd number. The common and difference parts of the conjugate precession angles $\mathbf{R}_{l,n}$ and $\mathbf{r}_{l,n}$ can be defined similarly to those for unit concatenation sequences. The precession vectors $\theta_{l,n}^{\pm} \equiv n\theta_l^{\pm}$ do not change their directions with increasing number of repeated units but only the angle of rotations grows linearly with n . Therefore,

$$r_{l,n}/R_{l,n} = r_l/R_l \cong r_{l-1}. \quad (88)$$

Since $R_{l,n} = |\sin \theta_{l,n}^+ / 2 + \sin \theta_{l,n}^- / 2| / 2 \leq 1$, the distinguishability between conjugate pseudo-spins given by equation (71) is bounded from above by

$$\delta_{l,n}^2 \leq 4r_{l,n}^2 \leq 4(r_l/R_l)^2 \cong 4r_{l-1}^2. \quad (89)$$

Since by each order of concatenation until the threshold level is reached, the common precession angle R_l is increased by a factor two and the difference part r_l is suppressed by a factor of $R_l \ll 1$, the ratio r_l/R_l can be made arbitrarily small. In the mesoscopic system, the number of excitation modes (pseudo-spins) is finite, so the coherence under repeated concatenation control is bounded from below by

$$\mathcal{L}_{+,-}(n\tau_l) \geq \exp \left(-2 \sum_k r_l^2 / R_l^2 \right) \cong \mathcal{L}_{+,-}(\tau_{l-1}). \quad (90)$$

The summation $\sum_k 2r_l^2 / R_l^2$ can be made arbitrarily small by choosing a small delay time and a sufficiently deep concatenation level, which means the electron spin coherence is stabilized virtually without decoherence. The coherence after n units of l th order concatenation sequences is no less than that after a single unit of $(l-1)$ th order concatenation sequence for $l > 1$. The analysis above is verified by numerical simulations presented in figure 18.

7. Summary and discussions

We have studied the electron spin decoherence in a mesoscopic bath of nuclear spins in semiconductor QDs. For QDs of interest in experiments, the electron–nuclear hyperfine interaction is much stronger than the nuclear spin interactions, so the electron spin and those nuclear spins in direct contact with the electron can be well isolated from the environment as a closed quantum system in the timescale of interest, giving the definition of a mesoscopic system.

When the longitudinal spin relaxation is absent (suppressed here by a moderate to strong magnetic field $\gtrsim 0.1$ T and a low temperature $\lesssim 1$ K), pure dephasing (or transverse relaxation) induced by the nuclear spin bath is the dominant cause for the loss of electron quantum coherence. In the quantum theory, decoherence is a consequence of the entanglement between the quantum object and the bath established by the evolution of bath along bifurcated pathways in the Hilbert space for different basis states of the quantum object. By contrast, semiclassical spectral diffusion theories ascribe the decoherence to accumulation of a random phase from a fluctuating local




Figure 18. The electron spin coherence under the control of repeated units of concatenated sequences, as functions of the number of units for a fixed pulse delay time $\tau = 0.2 \mu\text{s}$. The QD is under the same condition as in figure 17.

field. The nuclear spin dynamics, which is conditioned on the electron spin basis states, can be shepherded by manipulation of the electron spin. In particular, when the electron spin is flipped, the bifurcating nuclear spin pathways will exchange their evolution directions in the Hilbert space and could intersect at a later time, leading to the disentanglement of the electron from the nuclei and hence the restoration of the coherence even after it has been totally lost to the nuclear bath. Such recoherence by disentanglement is fundamentally different from spin echoes which result from re-focusing of random phases.

The solution to the many-body nuclear bath dynamics relies on four essential conditions. (i) The QD size is in the mesoscopic regime. (ii) The external field is much stronger than the hyperfine interaction. (iii) The temperature is low for the electron spin but high for the nuclear spins. (iv) The timescale considered is much shorter than the inverse nuclear interaction strength. Such conditions are realistic. They make possible two important simplifications in theory. Firstly, the electron spin dephasing in ensemble dynamics is factorized into two factors, namely, one due to the inhomogeneous broadening, and the other due to the entanglement between the electron spin and the nuclear spins. Secondly, the nuclear spin dynamics in the timescale of interest is approximated as independent excitations of pair-wise flip-flops. The solution of the bath quantum dynamics provides a simple geometrical basis for the design of pulse sequences for recoherence. The controllability of the mesoscopic bath dynamics, however, is a rather general concept independent of the simplifications made in the present paper.

A close examination of the single-system dynamics free of the inhomogeneous broadening reveals a wealth of phenomena in the electron spin decoherence. For example, a coherence recovery at a time different from the spin echo time would have been totally eclipsed in ensemble dynamics, since the dephasing due to inhomogeneous broadening is usually faster by orders of magnitude than the entanglement-induced decoherence. It is also shown that the single-pulse Hahn echo signals do not measure the electron spin decoherence in FID configuration: firstly, the decoherence caused by hyperfine-mediated nuclear pair-flips has a strong effect on the FID when the external field is not too large, but it is removed from the Hahn echo signal due to disentanglement; secondly, the decoherence by the intrinsic nuclear spin interaction is also partially suppressed, which results in a Hahn-echo decay time approximately $\sqrt{2}$ times the

FID time due to the intrinsic interaction. The decoherence behaviour in single-system dynamics (such as the recoherence at $\sqrt{2}\tau$) could eventually be directly observed in experiments if the inhomogeneous broadening can be filtered out, e.g. by a projective measurement of the local Overhauser field [20], [67]–[69].

To observe the recoherence by disentanglement in ensemble experiments [40, 71], pulse sequences are designed to force the disentanglement to coincide with a spin echo. The simplest solution is the Carr–Purcell sequence, which has one pulse at τ and another at 3τ and forces the disentanglement to take place at the echo time 4τ . More sophisticated designs such as concatenated sequences can be employed to suppress the decoherence to an arbitrary order of the pulse delay time in units of the inverse nuclear interaction strength, and a repetition of the concatenated sequences can preserve the electron spin coherence until other decoherence mechanisms (such as phonon scattering) become significant.

The design of concatenated pulse sequences for disentanglement resembles the dynamical decoupling schemes which have been developed in decades for NMR spectroscopies [48, 50, 51] and recently studied for quantum computation [54, 55, 57]. The disentanglement schemes exploit the controlling power of pulse sequences in a subtly different way from the dynamical decoupling schemes: the latter uses short pulses to flip the quantum object frequently so that the object–bath interaction is dynamically averaged to zero, while the former does not seek to eliminate the effective coupling but rather focuses on controlling the evolution of the bath wavefunction. For example, for local nuclear spin pair-flips in the single-pulse control configuration, the effective interaction $\hat{S}_e^z \otimes \hat{H}_{\text{eff}}$ at the disentanglement time $\sqrt{2}\tau$, defined by $\hat{H}_{\text{eff}} \equiv \hat{H}_{\text{eff}}^+ - \hat{H}_{\text{eff}}^-$ and $e^{-i\hat{H}_{\text{eff}}^{\pm}\sqrt{2}\tau} \equiv e^{-i\hat{H}^{\mp}(\sqrt{2}-1)\tau}e^{-i\hat{H}^{\pm}\tau}$, does not vanish even in the leading order of the hyperfine coupling. This is also the case for the disentanglement under the control of equally spaced pulses. One may question why a nonzero effective object–bath interaction could lead to vanishing decoherence. The reason is that the control steers the wavefunction evolution and the initial state of the bath is of some special form (namely, the nuclear spin states at the temperature under consideration possesses no off-diagonal coherence). The disentanglement time is in general dependent on the initial state (a universal disentanglement scheme independent of the initial state is only possible when the effective system–bath coupling vanishes). The dependence of the disentanglement on the initial state is implied in the control by evenly separated pulses shown in figure 14(a): for instance, if the nuclear spins present at $\sqrt{2}\tau$ are set as the initial state (which possesses off-diagonal two-spin coherence), the disentanglement after the electron spin flip at τ_0 would occur at $(\sqrt{3}-1)(\sqrt{2}+1)\tau_0$ (on the contrary, for initial states $|\mathcal{J}\rangle$ which has no off-diagonal coherence, the disentanglement occurs at $\sqrt{2}\tau_0$). It would be very interesting to study the decoherence and the disentanglement for spin baths with certain initial off-diagonal coherence such as in quantum memory applications [77].

Even though the concatenated disentanglement does cause the effective electron–nuclear coupling to vanish, the difference from the dynamical decoupling is still unambiguously evidenced by their different controlling small parameters. The controlling small parameter in dynamical decoupling is the object–bath interaction strength, and the pulse delay time should be much shorter than the inverse of that strength. In the dynamical disentanglement, the object–bath interaction strength is a quite irrelevant parameter, but the short-time condition is given by the bath interaction. Furthermore, in concatenated dynamical decoupling, the decoherence is eliminated in orders of the system–interaction strength times the pulse delay time, but in the disentanglement, the decoherence is suppressed in orders of the bath interaction strength times the delay time.

We conclude this paper by pointing out the limitations of the disentanglement scheme as an alternative to other methods of preserving coherence of quantum states such as dynamical decoupling. Firstly, the disentanglement method depends on the initial bath state, and hence the specific design of pulse sequences may vary from case to case (though to many commonly encountered cases such as thermalized baths the pulse sequences studied in this paper may be straightforwardly applied). Secondly, the disentanglement is possible only when the coherence of the mesoscopic system consisting of the quantum object and the bath is not lost to the macroscopic environment. Notwithstanding these limitations, the concept of disentanglement could be useful for small quantum systems, such as spins in molecules [77] and spins of nitrogen-vacancy centres in diamonds [78]. It should also be mentioned that even though the discussions in this paper have been largely facilitated by the pair-correlation approximation in treating the many-body nuclear spin dynamics, the disentanglement is a general notion independent of the employed approximation.

Acknowledgments

This work was supported by NSF DMR-0403406 and ARO/NSA-LPS.

Appendix A. Factorization of single-system dynamics and inhomogeneous broadening

The nuclear spin dynamics is basically determined by the excitation spectra of the pseudo-spins when the bath size is large enough. The excitation spectra for different nuclear spin configurations $|\mathcal{J}\rangle$ should be the same except for a relative variance in the order of $1/\sqrt{N}$ according to the central limit theorem in statistics. Here we give a numerical test of the single-system dynamics for a few initial nuclear spin configurations randomly selected from the thermal ensemble. The results are shown in figure A.1. Remarkably, the coherence for four initial states randomly selected from the thermal ensemble is visually indistinguishable, both in the FID regime ($t < 0.5 \mu\text{s}$) and after the controlling pulses are applied ($t > 0.5 \mu\text{s}$). The relative deviation between different curves is consistent with the estimate by the central limit theorem.

Appendix B. Boundary of mesoscopic nuclear spin bath

Intuitively, the nuclear spins in direct contact with the electron spin can be taken as the mesoscopic bath. The boundary of the bath is roughly defined by the condition that the hyperfine interaction is stronger than the nuclear spin interaction for nuclei within the bath, and otherwise for those without the bath. The interaction between the nuclear spins within and without the bath across the boundary, however, could make this definition somehow arbitrary. One can define a larger spin bath by including one or more layers of nuclei outside the boundary. Since the nuclear spin interaction is much weaker than the hyperfine interaction, the cross-boundary interaction should be unimportant in timescales of interest. Here, we present the numerical test of the dependence of the electron spin decoherence on the choice of the boundary of the mesoscopic nuclear spin bath, particularly by examining the real-time behaviour under the Carr–Purcell control. The result in figure B.1 confirms the assumption that the slight ambiguity in defining the mesoscopic bath as a closed system is unimportant.




Figure A.1. (a) The electron spin coherence under the control of a Carr–Purcell pulse sequence, for various initial nuclear spin states (indicated by arrows in figure 4): $|\mathcal{J}_1\rangle$ with local Overhauser field $E_{\mathcal{J}_1} = -0.5\Gamma_2^*$ (dot-dashed red line), $|\mathcal{J}_2\rangle$ with $E_{\mathcal{J}_2} = 0$ (solid black line), $|\mathcal{J}_3\rangle$ with $E_{\mathcal{J}_3} = 0.5\Gamma_2^*$ (dotted blue line), and $|\mathcal{J}_4\rangle$ with $E_{\mathcal{J}_4} = \Gamma_2^*$ (dashed green line). The four lines are indeed indistinguishable in the figure. (b) The deviation of the coherence for various initial states from that for the state $|\mathcal{J}_1\rangle$, $\Delta\mathcal{L}_{+,-}(t) \equiv |\langle\mathcal{J}_n|e^{i\hat{H}t}e^{-i\hat{H}_+t}|\mathcal{J}_n\rangle| - |\langle\mathcal{J}_1|e^{i\hat{H}t}e^{-i\hat{H}_+t}|\mathcal{J}_1\rangle|$, amplified by 1000 times. The InAs QD is of size $34 \times 34 \times 3 \text{ nm}^3$, at a temperature of 1 K, and under an external field of 10 tesla.

Appendix C. Pair-flip numbers

The number of pair-flips can be calculated with the pseudo-spin model, as an *a posteriori* check of the pair-correlation approximation. The general formalism is [4]

$$N_{\text{flip}}(t) \equiv \max_{\pm} \sum_k |\langle\uparrow|\psi_k^{\pm}(t)\rangle|^2 \approx \sum_k |\langle\uparrow|\psi_k^+(t)\rangle|^2. \quad (\text{C.1})$$

To be specific, let us consider the pair-flip number under concatenated control. With the notations defined in section 6, the pair-flip number under the l th order concatenation control is

$$N_{\text{flip}}^{(l)}(\tau_l) = \sum_k \left[(n_l^+)_x^2 + (n_l^+)_y^2 \right], \quad (\text{C.2})$$

where $(n_l^+)_x$ and $(n_l^+)_y$ are the x - and y -component of the precession vector \mathbf{n}_l^+ defined in equation (69), respectively, and $\tau_l \equiv 2^l\tau$. With the condition $\tau_l \ll B_k^{-1}$, the results in the leading order of the nuclear spin interaction strength are

$$N_{\text{flip}}^{(0)}(\tau) \cong N_{\text{flip},A}^{(0)}(\tau) + N_{\text{flip},B}^{(0)}(\tau), \quad (\text{C.3a})$$

$$N_{\text{flip},A}^{(0)}(\tau) \cong \sum_{k \in \mathbb{G}_A} A_k^2 \tau^2 \text{sinc}^2 \frac{E_k \tau}{2} \lesssim \frac{A_\alpha^4 \tau^2}{N^2 \Omega_c^2}, \quad (\text{C.3b})$$




Figure B.1. (a) The electron spin coherence under the control of a Carr–Purcell pulse sequence, for various choices of boundary of the nuclear spin bath: for the dashed red, solid black, and dotted blue lines, the boundary is set 0, 1 and 3 layers of unit cells outside the boundary defined by the QD potential walls. The three lines are indistinguishable in the figure. (b) The deviation (amplified by 100 times) of the coherence for various choices of the bath boundary, referenced to the first choice of boundary. The QD is as in figure A.1.

$$N_{\text{flip},B}^{(0)}(\tau) \cong \sum_{k \in \mathbb{G}_B} B_k^2 \tau^2 \text{sinc}^2 \frac{E_k \tau}{2} \lesssim N B_k^2 \tau^2, \quad (\text{C.3c})$$

$$N_{\text{flip}}^{(l>0)}(\tau_l) \cong \sum_{k \in \mathbb{G}_B} B_k^2 \tau_l^2 \text{sinc}^2(E_k \tau) \lesssim N B_k^2 \tau_l^2. \quad (\text{C.3d})$$

Here the number of local pair-flips and that for hyperfine-mediated nonlocal pair-flips in FID configuration are identified with the subscripts B and A , respectively. As illustrated in figure 11(b), the pseudo-spin precession for nonlocal pair-flips is reversed when the electron spin is flipped, so under the concatenated control the number of hyperfine-mediated pair-flips will oscillate periodically, with the maximum value $N_{\text{flip},A}^{(0)}(\tau)$ at $(2n+1)\tau$ and minimum value 0 at $2n\tau$. At the spin echo time τ_l for $l > 0$, the number of pair-flips is contributed by the intrinsic nuclear spin interaction. The number of pair-flips is plotted in figure C.1 for a typical QD in the Carr–Purcell control configuration, showing the features discussed above.

To justify the pair-correlation approximation, the small pair-flip number condition

$$N_{\text{flip}}^2 \ll N, \quad (\text{C.4})$$

is to be fulfilled (see appendix D). Thus, the criteria for the timescales τ and τ_l being short are given by

$$N B_k^4 \tau_l^4 \ll 1, \quad (\text{C.5a})$$

$$N^{-5} \Omega_e^{-4} \mathcal{A}_\alpha^8 \tau^4 \ll 1, \quad (\text{C.5b})$$




Figure C.1. (a) The electron spin coherence in the Carr–Purcell control configuration. (b) The number of pair-flips due to the intrinsic nuclear spin interaction (dotted red line) and that due to the hyperfine-mediated interaction (solid blue line, amplified by 10 times). The InAs QD is as in figure A.1 with $B = 2$ tesla.

for consideration of the intrinsic interaction and the hyperfine-mediated interaction, respectively. For an InAs QD containing about 10^5 nuclei under an external field of $B_{\text{ext}} \sim 1$ tesla, the valid timescales is estimated to be

$$\tau_l \lesssim 100 \mu\text{s}, \quad (\text{C.6a})$$

$$\tau \lesssim 10 \mu\text{s}. \quad (\text{C.6b})$$

A natural timescale of interest in experiments is the decay time of the Hahn echo signal, which is estimated from equation (43b) and (50) to be

$$T_{\text{H},B} \sim N^{5/12} B_k^{-1/2} \mathcal{A}_\alpha^{-1/2}. \quad (\text{C.7})$$

This timescale together with equation (C.5) sets ranges for the field strength and the QD size where the pair-correlation approximation is valid:

$$N^{8/3} B_k^2 \mathcal{A}_\alpha^{-2} \ll 1, \quad (\text{C.8a})$$

$$N^{-10/3} \Omega_e^{-4} B_k^{-2} \mathcal{A}_\alpha^6 \ll 1, \quad (\text{C.8b})$$

as schematically shown in figure C.1.

Appendix D. Error estimation for pair-correlation approximation

To estimate the error of the pair-correlation approximation, we express the exact and the approximate nuclear spin wavefunctions as

$$|\mathcal{J}_{\text{exact}}^\pm(t)\rangle = |\mathcal{J}_{\text{exact,uncorr}}^\pm\rangle + |\mathcal{J}_{\text{exact,corr}}^\pm\rangle, \quad (\text{D.1a})$$




Figure C.2. Schematics for the range of the QD size and the field strength (shaded area) where the pair-correlation approximation is justified in the timescales of interest in the single-pulse Hahn echo configuration. The upper and lower bounds set limits on the number of the intrinsic local pair-flips and on that of the hyperfine-mediated nonlocal pair-flips, respectively.

$$|\mathcal{J}_{\text{PCA}}^{\pm}(t)\rangle = |\mathcal{J}_{\text{PCA,uncorr}}^{\pm}\rangle + |\mathcal{J}_{\text{PCA,corr}}^{\pm}\rangle, \quad (\text{D.1b})$$

respectively, where $|\mathcal{J}_{\text{exact,corr}}^{\pm}\rangle$ and $|\mathcal{J}_{\text{PCA,corr}}^{\pm}\rangle$ denote the wavefunctions containing correlated pair-flips, and $|\mathcal{J}_{\text{exact,uncorr}}^{\pm}\rangle$ and $|\mathcal{J}_{\text{PCA,uncorr}}^{\pm}\rangle$ are the parts containing uncorrelated pair-flips. The evolution starts with a randomly chosen nuclear spin state $|\mathcal{J}\rangle$. The uncorrelated part of the wavefunction in the exact solution and that in the approximation are determined by the same Green's function $\hat{G}_0(t, t')$ and by the probability amplitudes at the initial state $C_{\text{PCA},\mathcal{J}}^{\pm}(t')$ and $C_{\text{exact},\mathcal{J}}^{\pm}(t')$ as

$$|\mathcal{J}_{\text{exact,uncorr}}^{\pm}\rangle = \int \hat{G}_0(t, t') C_{\text{exact},\mathcal{J}}^{\pm}(t') |\mathcal{J}\rangle dt', \quad (\text{D.2a})$$

$$|\mathcal{J}_{\text{PCA,uncorr}}^{\pm}\rangle = \int \hat{G}_0(t, t') C_{\text{PCA},\mathcal{J}}^{\pm}(t') |\mathcal{J}\rangle dt'. \quad (\text{D.2b})$$

After a few pair-flips have occurred while still well before the correlated part of the wavefunction becomes significant, the probability amplitude of the initial state $|\mathcal{J}\rangle$ decays already to zero. So the amplitudes at the initial state are essentially the same in the exact and the approximate solutions. Consequently, the wavefunctions containing only uncorrelated pair-flips are equal in the two solutions

$$|\mathcal{J}_{\text{exact,uncorr}}^{\pm}\rangle = |\mathcal{J}_{\text{PCA,uncorr}}^{\pm}\rangle. \quad (\text{D.3a})$$

Thus the error of the pair-correlation approximation in calculating the electron spin coherence correlation is estimated to be

$$\delta\mathcal{L}_{+,-}^s \sim |\langle \mathcal{J}_{\text{exact,corr}}^{-} | \mathcal{J}_{\text{exact,corr}}^{+} \rangle - \langle \mathcal{J}_{\text{PCA,corr}}^{-} | \mathcal{J}_{\text{PCA,corr}}^{+} \rangle| \leq P_{\text{corr}}, \quad (\text{D.4})$$

where P_{corr} is the probability of having correlated pair-flips. When the number of pair-flips $N_{\text{flip}} \ll N$, $P_{\text{corr}} \sim 1 - \exp(-qN_{\text{flip}}^2/N)$ (q is a factor close to the number of nearest neighbours of a nuclear spin).

References

- [1] Zurek W H 2003 *Rev. Mod. Phys.* **75** 715
- [2] Joos E, Zeh H D, Kiefer C, Giulini D, Kupsch J and Stamatescu I-O 2003 *Decoherence and the Appearance of a Classical World in Quantum Theory* (New York: Springer)
- [3] Schlosshauer M 2004 *Rev. Mod. Phys.* **76** 1267
- [4] Yao W, Liu R B and Sham L J 2006 *Phys. Rev. B* **74** 195301
- [5] Yao W, Liu R B and Sham L J 2007 *Phys. Rev. Lett.* **98** 077602
- [6] Saikin S K, Yao W and Sham L J 2007 *Phys. Rev. B* **75** 125314
- [7] Witzel W M, de Sousa R and Das Sarma S 2005 *Phys. Rev. B* **72** 161306(R)
- [8] Witzel W M and Das Sarma S 2007 *Phys. Rev. Lett.* **98** 077601
- [9] Witzel W M and Das Sarma S 2006 *Phys. Rev. B* **74** 035322
- [10] Anderson P W 1954 *J. Phys. Soc. Japan* **9** 316
- [11] Schulten K and Wolynes P G 1978 *J. Chem. Phys.* **68** 3292
- [12] Imry Y 1997 *Introduction to Mesoscopic Physics* (New York: Oxford University Press)
- [13] Leggett A, Chakravarty S, Dorsey A, Fisher M, Garg A and Zwerger W 1987 *Rev. Mod. Phys.* **59** 1
- [14] Loss D and DiVincenzo D P 1998 *Phys. Rev. A* **57** 120
- [15] Imamoglu A, Awschalom D D, Burkard G, DiVincenzo D P, Loss D, Sherwin M and Small A 1999 *Phys. Rev. Lett.* **83** 4204
- [16] Awschalom D D, Loss D and Samarth N (ed) 2002 *Semiconductor Spintronics and Quantum Computation* (New York: Springer)
- [17] Merkulov I A, Efros A L and Rosen M 2002 *Phys. Rev. B* **65** 205309
- [18] Khaetskii A V, Loss D and Glazman L 2002 *Phys. Rev. Lett.* **88** 186802
- [19] Khaetskii A, Loss D and Glazman L 2003 *Phys. Rev. B* **67** 195329
- [20] Coish W A and Loss D 2004 *Phys. Rev. B* **70** 195340
- [21] Semenov Y G and Kim K W 2003 *Phys. Rev. B* **67** 073301
- [22] Deng C and Hu X 2006 *Phys. Rev. B* **73** 241303(R)
- [23] Shenvi N, de Sousa R and Whaley K B 2005 *Phys. Rev. B* **71** 224411
- [24] Shenvi N, de Sousa R and Whaley K B 2005 *Phys. Rev. B* **71** 144419
- [25] de Sousa R and Das Sarma S 2003 *Phys. Rev. B* **68** 115322
- [26] de Sousa R and Das Sarma S 2003 *Phys. Rev. B* **67** 033301
- [27] Khaetskii A V and Nazarov Y 2000 *Phys. Rev. B* **61** 12639
- [28] Woods L M, Reinecke T L and Lyanda-Geller Y 2002 *Phys. Rev. B* **66** 161318
- [29] Golovach V N, Khaetskii A and Loss D 2004 *Phys. Rev. Lett.* **93** 016601
- [30] Semenov Y G and Kim K W 2004 *Phys. Rev. Lett.* **92** 026601
- [31] Fujisawa T, Austing D G, Tokura Y, Hirayama Y and Tarucha S 2002 *Nature* **419** 278
- [32] Elzerman J M, Hanson R, Willems van Beveren L H, Witkamp B, Vandersypen L M K and Kouwenhoven L P 2004 *Nature* **430** 431
- [33] Kroutvar M, Ducommun Y, Heiss D, Bichler M, Schuh D, Abstreiter D and Finley J J 2004 *Nature* **432** 81
- [34] Johnson A C, Petta J R, Taylor J M, Yacoby A, Lukin M D, Marcus C M, Hanson M P and Gossard A C 2005 *Nature* **435** 925
- [35] Braun P *et al* 2005 *Phys. Rev. Lett.* **94** 116601
- [36] Atatüre M, Dreiser J, Högele A, Karrai K and Imamoglu A 2006 *Science* **312** 551
- [37] Gurudev Dutt M V *et al* 2005 *Phys. Rev. Lett.* **94** 227403
- [38] Bracker A S *et al* 2005 *Phys. Rev. Lett.* **94** 047402
- [39] Koppens F H L, Folk J A, Elzerman J M, Hanson R, van Beveren L H W, Vink I T, Tranitz H P, Wegscheider W, Kouwenhoven L P and Vandersypen L M K 2005 *Science* **309** 1346
- [40] Petta J R, Johnson A C, Taylor J M, Laird E A, Yacoby A, Lukin M D, Marcus C M, Hanson M P and Gossard A C 2005 *Science* **309** 2180

- [41] Greilich A, Yakovlev D R, Shabaev A, Efros A L, Yugova I A, Oulton R, Stavarache V, Reuter D, Wieck A and Bayer M 2006 *Science* **313** 341
- [42] Prokof'ev N V and Stamp P C E 2000 *Rep. Prog. Phys.* **63** 669
- [43] Schliemann J, Khaetskii A V and Loss D 2002 *Phys. Rev. B* **66** 245303
- [44] de Sousa R, Shenvi N and Whaley K B 2005 *Phys. Rev. B* **72** 045310
- [45] Brune M, Hagley E, Dreyer J, Maitre X, Maali A, Wunderlich C, Raimond J M and Haroche S 1996 *Phys. Rev. Lett.* **77** 4887
- [46] Griffiths R B 2002 *Consistent Quantum Theory* (New York: Cambridge University Press)
- [47] Taylor J M, Imamoglu A and Lukin M D 2003 *Phys. Rev. Lett.* **91** 246802
- [48] Rhim W, Pines A and Waugh J 1970 *Phys. Rev. Lett.* **25** 218
- [49] Zhang S, Meier B H and Ernst R R 1992 *Phys. Rev. Lett.* **69** 2149
- [50] Haeberle U and Waugh J 1968 *Phys. Rev.* **175** 453
- [51] Slichter C P 1992 *Principles of Magnetic Resonance* 3rd edn (Berlin: Springer-Verlag)
- [52] Viola L and Lloyd S 1998 *Phys. Rev. A* **58** 2733
- [53] Viola L, Knill E and Lloyd S 1999 *Phys. Rev. Lett.* **82** 2417
- [54] Viola L and Knill E 2005 *Phys. Rev. Lett.* **94** 060502
- [55] Khodjasteh K and Lidar D A 2005 *Phys. Rev. Lett.* **95** 180501
- [56] Khodjasteh K and Lidar D A 2006 *Preprint quant-ph/0607086*
- [57] Kern O and Alber G 2005 *Phys. Rev. Lett.* **95** 250501
- [58] Jacobi K 2003 *Prog. Surf. Sci.* **71** 185
- [59] Bloembergen N and Rowland T J 1955 *Phys. Rev.* **97** 1679
- [60] Sundfors R K 1969 *Phys. Rev.* **185** 458
- [61] Paget D, Lampel G and Sapoval B 1977 *Phys. Rev. B* **15** 5780
- [62] Cucchiatti F M, Dalvit D A R, Paz J P and Zurek W H 2003 *Phys. Rev. Lett.* **91** 210403
- [63] Cucchiatti F M, Paz J P and Zurek W H 2005 *Phys. Rev. A* **72** 052113
- [64] Quan H T, Song Z, Liu X F, Zanardi P and Sun C P 2006 *Phys. Rev. Lett.* **96** 140604
- [65] Gupta J A, Awschalom D D, Efros A L and Rodina A V 2002 *Phys. Rev. B* **66** 125307
- [66] Koppens F H L, Buizert C, Tielrooij K J, Vink I T, Nowack K C, Meunier T, Kouwenhoven L P and Vandersypen L M K 2006 *Nature* **442** 766
- [67] Giedke G, Taylor J M, D'Alessandro D, Lukin M D and Imamoglu A 2005 *Preprint quant-ph/0508144*
- [68] Klauser D, Coish W A and Loss D 2006 *Phys. Rev. B* **73** 205302
- [69] Stepanenko D, Burkard G, Giedke G and Imamoglu A 2006 *Phys. Rev. Lett.* **96** 136401
- [70] Liu R B, Yao W and Sham L J 2005 *Phys. Rev. B* **72** 081306(R)
- [71] Tyryshkin A M, Lyon S A, Astashkin A V and Raitsimring A M 2003 *Phys. Rev. B* **68** 193207
- [72] Abe E, Itoh K M, Isoya J and Yamasaki S 2004 *Phys. Rev. B* **70** 033204
- [73] Abe E, Isoya J and Itoh K M 2005 *J. Supercond.* **18** 157
- [74] Chen P, Piermarocchi C, Sham L J, Gammon D and Steel D G 2003 *Phys. Rev. B* **69** 075320
- [75] Chen S W and Liu R B unpublished
- [76] Taylor J M, Marcus C M and Lukin M D 2003 *Phys. Rev. Lett.* **90** 206803
- [77] Lehmann J, Gaita-Ariño A, Coronado E and Loss D 2007 *Nat. Nanotechnol.* **2** 312
- [78] Gurudev Dutt M V, Childress L, Jiang L, Togan E, Maze J, Jelezko F, Zibrov A S, Hemmer P R and Lukin M D 2007 *Science* **316** 1312