



Heavy Fermions and symplectic symmetry

R. Flint¹


M. Dzero^{1,2}

P. Coleman¹

¹ Rutgers U.

² Columbia U.

[arXiv:0710.1126](https://arxiv.org/abs/0710.1126)





Heavy Fermions and symplectic symmetry

R. Flint¹

M. Dzero^{1,2}


P. Coleman¹

¹ Rutgers U.


² Columbia U.

[arXiv:0710.1126](https://arxiv.org/abs/0710.1126)

- Collision of ideas
- New Superconductors
- Time reversal and Symplectic spins
- Super-coherent Kondo
- Predictions




Heavy Fermions: Collision of ideas




Tuson Park, (2007).

CeRhIn₅




Heavy Fermions: Collision of ideas




Tuson Park, (2007).


CeRhIn₅




Heavy Fermions: Collision of ideas




Custers et al (2003)




Heavy Fermions: Collision of ideas



Heavy Fermions: Collision of ideas




Heavy Fermions: Collision of ideas




| Oxides | | Intermetallics | |
|---|-------|----------------|-------------------------------|
| | T_c | T_c | |
| <u>?</u> | | 18.5K | <u>PuCoGa₅</u> '02 |
| <u>YBa₂Cu₃O₇</u> | 92K | 2K | <u>CeCoIn₅</u> '01 |
| <u>Ba2212</u> | 12K | 0.2K | <u>CeIn₃</u> '96 |

Heavy Fermions: Collision of ideas



“Fruit fly” for correlated materials

Heavy Fermions: Collision of ideas




| Oxides | | Intermetallics | |
|---|-------|----------------|---|
| | T_c | T_c | |
| ? | | 18.5K | <u>PuCoGa₅</u> '02 |
| | | 4.5K | <u>NpPd₂Al₅</u> '07 |
| <u>YBa₂Cu₃O₇</u> | 92K | 2K | <u>CeCoIn₅</u> '01 |
| <u>Ba2212</u> | 12K | 0.2K | <u>CeIn₃</u> '96 |

"Fruit fly" for correlated materials

New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5


J. Sarrao et al., Nature 420, (2002)



18.5 K Heavy Fermion
SC PuCoGa_5


New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5

J. Sarrao et al., Nature 420, (2002)



18.5 K Heavy Fermion
SC PuCoGa_5


D. Aoki et al., JPSJ 76 (2007)




4.5K Heavy Fermion S.C
 NpPd_2Al_5

New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5

J. Sarrao et al., Nature 420, (2002)




18K Heavy Fermion S. C
 PuCoGa_5




New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5

J. Sarrao et al., Nature 420, (2002)




18K Heavy Fermion S. C
 PuCoGa_5



Directly Curie PM to SC


New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5

D. Aoki et al., JPSJ 76 (2007)




4.5K Heavy Fermion S.C
 NpAl_2Pd_5

New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5




Directly Curie PM to SC

D. Aoki et al., JPSJ 76 (2007)




4.5K Heavy Fermion S.C
 NpAl_2Pd_5

New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5




Directly Curie PM to SC

D. Aoki et al., JPSJ 76 (2007)




4.5K Heavy Fermion S.C
 NpAl_2Pd_5




Substantial fraction of
spin entropy.
 $\frac{1}{3} R \ln 2$


New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5




Directly Curie PM to SC




Substantial fraction of
spin entropy.
 $\frac{1}{3} R \ln 2$




New HF Superconductors: PuCoGa_5 & NpAl_2Pd_5



Directly Curie PM to SC



Substantial fraction of
spin entropy.
 $\frac{1}{3} R \ln 2$



- Suggests Kondo spin quenching and superconductivity develop simultaneously.
- Spin is not the glue, but the *fabric*

Symplectic Spins and Large N Expansions

Strongly correlated electron physics: no small parameter

Strongly correlated electron physics: no small parameter


Large N : family of models which retain the key physics and can be solved in the large N limit.

$$Z = \sum_{configs} e^{-N \times S[\psi]}$$

Strongly correlated electron physics: no small parameter

Large N : family of models which retain the key physics and can be solved in the large N limit.



$$Z = \sum_{configs} e^{-N \times S[\psi]}$$



Strongly correlated electron physics: no small parameter

Large N : family of models which retain the key physics and can be solved in the large N limit.

$$Z = \sum_{configs} e^{-N \times S[\psi]}$$



The magnetism and neutrality of the electron spin are manifested two discrete parities:

odd under time reversal,
even under charge conjugation

The magnetism and neutrality of the electron spin are manifested two discrete parities:

odd under time reversal,
even under charge conjugation

$$\begin{array}{ll} S \xrightarrow{\theta} -S & \text{Magnetism} \\ S \xrightarrow{C} +S & \text{Neutrality} \end{array}$$

The magnetism and neutrality of the electron spin are manifested two discrete parities:

odd under time reversal,
even under charge conjugation

$$\begin{array}{ll} S \xrightarrow{\theta} -S & \text{Magnetism} \\ S \xrightarrow{C} +S & \text{Neutrality} \end{array}$$

These discrete parities are lost in conventional large N expansions.


Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K \quad (K a |\psi\rangle = a^* K |\psi\rangle)$$

Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$


$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$


$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$


$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$



$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$



$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$



$$(K a |\psi\rangle = a^* K |\psi\rangle)$$



Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K$$

$$(K a |\psi\rangle = a^* K |\psi\rangle)$$





Time reversal = antiunitary operator = $U K$

$$\theta = i\sigma_2 K \quad (K a|\psi\rangle = a^* K|\psi\rangle)$$

Invariance of time reversal under rotations U :

$$U\theta = \theta U \quad (U = e^{i\frac{\vec{\alpha}}{2} \cdot \vec{\sigma}})$$



Invariance of time reversal under rotations U:



$$\theta = i\sigma_2 K$$

$$U\theta = \theta U$$

$$(U = e^{i\frac{\vec{\alpha}}{2} \cdot \vec{\sigma}})$$

So that:

$$U\theta U^\dagger = \theta$$



Invariance of time reversal under rotations U:

$$\theta = i\sigma_2 K$$



$$U\theta = \theta U$$

$$(U = e^{i\frac{\vec{\sigma}}{2} \cdot \vec{\sigma}})$$

So that:

$$U\theta U^\dagger = \theta \quad \text{but}$$

$$KU^\dagger = (U^\dagger)^* K = U^T K$$



Invariance of time reversal under rotations U:

$$\theta = i\sigma_2 K$$

$$U\theta = \theta U$$

$$(U = e^{i\frac{\vec{\alpha}}{2} \cdot \vec{\sigma}})$$



So that:

$$U\theta U^\dagger = \theta \quad \text{but}$$

$$KU^\dagger = (U^\dagger)^* K = U^T K$$

$$U i\sigma_2 U^T = i\sigma_2$$

SYMPLECTIC CONDITION



SU(N) spin:

$$M_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta}$$

$$(\tilde{\alpha} \equiv \text{sgn}(\alpha))$$

SU(N) spin:

$$M_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta}$$

$$\xrightarrow{\theta} \tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$$\xrightarrow{c} \tilde{\alpha} \tilde{\beta} f_{-\beta} f_{-\alpha}^{\dagger} = -\tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$$(\tilde{\alpha} \equiv \text{sgn}(\alpha))$$

SU(N) spin:

$$M_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta}$$

$$\xrightarrow{\theta} \tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$$\xrightarrow{\mathcal{C}} \tilde{\alpha} \tilde{\beta} f_{-\beta} f_{-\alpha}^{\dagger} = -\tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$\mathcal{C}\theta = -1$, but \mathcal{C} and θ ill-defined.

$$(\tilde{\alpha} \equiv \text{sgn}(\alpha))$$

SU(N) spin:

$$M_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta}$$

$$\xrightarrow{\theta} \tilde{\alpha}\tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$$\xrightarrow{\mathcal{C}} \tilde{\alpha}\tilde{\beta} f_{-\beta} f_{-\alpha}^{\dagger} = -\tilde{\alpha}\tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$\mathcal{C}\theta = -1$, but \mathcal{C} and θ ill-defined.

“magnetic” moments

$$S_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta} - \tilde{\alpha}\tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}, \quad (\theta, \mathcal{C}) = (-, +)$$

“electric” dipoles

$$\mathcal{P}_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta} + \tilde{\alpha}\tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}, \quad (\theta, \mathcal{C}) = (+, -)$$

$S_{\alpha\beta} \equiv$ generators of the SP(N) group

$$(\tilde{\alpha} \equiv \text{sgn}(\alpha))$$

SU(N) spin:

$$M_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta}$$

$$\xrightarrow{\theta} \tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$$\xrightarrow{\mathcal{C}} \tilde{\alpha} \tilde{\beta} f_{-\beta} f_{-\alpha}^{\dagger} = -\tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}$$

$\mathcal{C}\theta = -1$, but \mathcal{C} and θ ill-defined.

“magnetic” moments

$$S_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta} - \tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}, \quad (\theta, \mathcal{C}) = (-, +)$$

“electric” dipoles

$$\mathcal{P}_{\alpha\beta} = f_{\alpha}^{\dagger} f_{\beta} + \tilde{\alpha} \tilde{\beta} f_{-\beta}^{\dagger} f_{-\alpha}, \quad (\theta, \mathcal{C}) = (+, -)$$

$S_{\alpha\beta} \equiv$ generators of the SP(N) group


$$[S, \tilde{\alpha} f_{\alpha} f_{-\alpha}] = 0$$

Singlet pair commutes with symplectic spin


→ Local SU(2) gauge symmetry.

Application to the Kondo Lattice

SPIN NEUTRALITY (Flint, Dzero,PC).




SPIN NEUTRALITY (Flint, Dzero, PC).



$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

SPIN NEUTRALITY (Flint, Dzero, PC).




$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

What is the gauge
field of the
constraint?

(Read Newns, PC, Mattis Lee... 80's)

SPIN NEUTRALITY (Flint, Dzero,PC).



$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$


$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)

What is the gauge field of the constraint?

(Read Newns, PC, Mattis Lee... 80's)

SPIN NEUTRALITY (Flint, Dzero,PC).



$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)


$$\begin{aligned} f_\alpha &\rightarrow e^{i\theta} f_\alpha \\ f_\alpha &\rightarrow \cos \phi f_\alpha + \sin \phi \text{sgn}(\alpha) f_{-\alpha}^\dagger \end{aligned}$$

SU(2) gauge symmetry (Affleck et al '89)

What is the gauge field of the constraint?

(Read Newns, PC, Millis Lee... 80's)

SPIN NEUTRALITY (Flint, Dzero,PC).



$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)


$$H_g = (\Phi - \mu) c_j^\dagger c_j + W_+(f^\dagger f^\dagger) + W_0(f^\dagger f) + W_-(ff)$$

$$U(1)_{\text{global}} \times SU(2)_{\text{local}}$$

What is the gauge field of the constraint?

(Read Newns, PC, Millis Lee... 80's)

SPIN NEUTRALITY (Flint, Dzero,PC).



What is the gauge field of the constraint?
(Read Newns, PC, Millis Lee... 80's)

$$H = \sum_k \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$


$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)

$$H_g = (\Phi - \mu) c_j^\dagger c_j + W_+(f^\dagger f^\dagger) + W_0(f^\dagger f) + W_-(ff)$$

$$U(1)_{\text{global}} \times SU(2)_{\text{local}}$$

SPIN NEUTRALITY (Flint, Dzero,PC).



What is the gauge field of the constraint?
(Read Newns, PC, Millis Lee... 80's)

$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)

$$H_g = (\Phi - \mu) c_j^\dagger c_j + W_+(f^\dagger f^\dagger) + W_0(f^\dagger f) + W_-(ff)$$


$$U(1)_{\text{global}} \times SU(2)_{\text{local}}$$

Φ

W^- W^0 W^+

cf Weinberg
Salem

SPIN NEUTRALITY (Flint, Dzero,PC).



What is the gauge field of the constraint?
(Read Newns, PC, Millis Lee... 80's)


$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$

$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$


"Symplectic spin" (Flint,Dzero,PC)

$$H_g = (\Phi - \mu) c_j^\dagger c_j + W_+(f^\dagger f^\dagger) + W_0(f^\dagger f) + W_-(f f)$$


$$U(1)_{\text{global}} \times SU(2)_{\text{local}}$$



SPIN NEUTRALITY (Flint, Dzero,PC).



What is the gauge field of the constraint?
(Read Newns, PC, Millis Lee... 80's)



$$H = \sum \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \frac{J}{N} \sum_j c_{j\alpha}^\dagger c_{j\beta} S_{\beta\alpha}(j) + H_g$$


$$S_{\alpha\beta} = f_\alpha^\dagger f_\beta - \text{sgn}(\alpha)\text{sgn}(\beta) f_{-\beta}^\dagger f_{-\alpha}$$

"Symplectic spin" (Flint,Dzero,PC)


$$H_g = (\Phi - \mu) c_j^\dagger c_j + W_+(f^\dagger f^\dagger) + W_0(f^\dagger f) + W_-(f f)$$

$$U(1)_{\text{global}} \times SU(2)_{\text{local}}$$

$$\text{photon} = (\Phi + W_0)/2$$
$$Z = (\Phi - W_0)/2$$



Model for PuCoGa_5 & NpPd_2Al_5




Model for PuCoGa₅ & NpPd₂Al₅

Tetragonal CF

| | n |
|-----|---|
| Pu: | 5 |
| Np: | 3 |

Pu moment

f^5




f^{n+1}

Γ_2

f^{n-1}

Γ_1

f^n




Model for PuCoGa₅ & NpPd₂Al₅

Tetragonal CF

| | n |
|-----|---|
| Pu: | 5 |
| Np: | 3 |

Pu moment

f^5





Virtual valence fluctuations mediate
Kondo-spin exchange in two channels with
different symmetry (Cox, 89).

Model for PuCoGa₅ & NpPd₂Al₅

Tetragonal CF

| | n |
|-----|---|
| Pu: | 5 |
| Np: | 3 |



Virtual valence fluctuations mediate Kondo-spin exchange in two channels with different symmetry (Cox, 89).

$$H = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + \frac{1}{N} \sum_{\mathbf{k}, \mathbf{k}'} \left(J_1 \psi_{1a}^{\dagger}(j) \psi_{1b}(j) + J_2 \psi_{2a}^{\dagger}(j) \psi_{2b}(j) \right) S^{ba}(j)$$

Single FS, two channels.

$$\psi_{\Gamma}(j) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \gamma_{\Gamma \mathbf{k}} c_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{x}_j}$$

Decoupling the interactions

$$\begin{aligned} H_I &= \frac{J_K}{N} \sum \psi_a^\dagger \psi_b \overbrace{\left(f_b^\dagger f_a - \tilde{a} \tilde{b} f_{-a}^\dagger f_{-b} \right)}^{S_{ba}} \\ &= -\frac{J_K}{N} \sum \left[(\psi^\dagger f)(f^\dagger \psi) + (\psi^\dagger \sigma_2 f^\dagger)(f \sigma_2 \psi) \right]. \end{aligned}$$

Decoupling the interactions

$$\begin{aligned}
 H_I &= \frac{J_K}{N} \sum \psi_a^\dagger \psi_b \overbrace{\left(f_b^\dagger f_a - \tilde{a} \tilde{b} f_{-a}^\dagger f_{-b} \right)}^{S_{ba}} \\
 &= -\frac{J_K}{N} \sum \left[(\psi^\dagger f)(f^\dagger \psi) + (\psi^\dagger \sigma_2 f^\dagger)(f \sigma_2 \psi) \right].
 \end{aligned}$$

$$H_I \rightarrow \left[\underbrace{[V f^\dagger + (\Delta f \sigma_2)] \psi + \text{H.c}}_{= \tilde{f}^\dagger \sqrt{|V|^2 + |\Delta|^2} = \tilde{f}^\dagger \tilde{V}} \right] + N \left(\frac{\bar{V} V + \bar{\Delta} \Delta}{J_K} \right).$$

- “Pairing” can be gauged away for 1 channel.

Decoupling the interactions

$$\begin{aligned}
 H_I &= \frac{J_K}{N} \sum \psi_a^\dagger \psi_b \overbrace{\left(f_b^\dagger f_a - \tilde{a} \tilde{b} f_{-a}^\dagger f_{-b} \right)}^{S_{ba}} \\
 &= -\frac{J_K}{N} \sum \left[(\psi^\dagger f)(f^\dagger \psi) + (\psi^\dagger \sigma_2 f^\dagger)(f \sigma_2 \psi) \right].
 \end{aligned}$$

$$H_I \rightarrow \left[[\tilde{V} \tilde{f}^\dagger] \psi + \text{H.c} \right] + N \left(\frac{\tilde{V}^2}{J_K} \right).$$

- “Pairing” can be gauged away for 1 channel.

Decoupling the interactions

- Two channels:

Decoupling the interactions

- Two channels:

$$H_I \rightarrow \left[[V_1 f^\dagger + (\Delta_1 f \sigma_2)] \psi_1 + \text{H.c} \right] + N \left(\frac{V_1 V_1 + \Delta_1 \Delta_1}{J_1} \right) \\ + \left[[V_2 f^\dagger + (\Delta_2 f \sigma_2)] \psi_2 + \text{H.c} \right] + N \left(\frac{\bar{V}_2 V_2 + \bar{\Delta}_2 \Delta_2}{J_2} \right)$$

Decoupling the interactions

- Two channels:

$$H_I \rightarrow \left[[V_1 f^\dagger + (\Delta_1 f \sigma_2)] \psi_1 + \text{H.c} \right] + N \left(\frac{V_1 V_1 + \Delta_1 \Delta_1}{J_1} \right) \\ + \left[[V_2 f^\dagger + (\Delta_2 f \sigma_2)] \psi_2 + \text{H.c} \right] + N \left(\frac{\bar{V}_2 V_2 + \bar{\Delta}_2 \Delta_2}{J_2} \right)$$

Gauge fixing does not remove the pairing

$$\rightarrow \left[[V_1 f^\dagger \psi_1 + (\Delta_2 f \sigma_2) \psi_2] + \text{H.c} \right] + N \left(\frac{V_1 V_1}{J_1} + \frac{\Delta_2 \Delta_2}{J_2} \right)$$

Decoupling the interactions

- Two channels:

$$H_I \rightarrow \left[[V_1 f^\dagger + (\Delta_1 f \sigma_2)] \psi_1 + \text{H.c} \right] + N \left(\frac{V_1 V_1 + \Delta_1 \Delta_1}{J_1} \right) \\ + \left[[V_2 f^\dagger + (\Delta_2 f \sigma_2)] \psi_2 + \text{H.c} \right] + N \left(\frac{\bar{V}_2 V_2 + \bar{\Delta}_2 \Delta_2}{J_2} \right)$$

Gauge fixing does not remove the pairing

$$\rightarrow \left[[V_1 f^\dagger \psi_1 + (\Delta_2 f \sigma_2) \psi_2] + \text{H.c} \right] + N \left(\frac{V_1 V_1}{J_1} + \frac{\Delta_2 \Delta_2}{J_2} \right)$$

The gauge-invariant cross term $V_1 \Delta_2 - V_2 \Delta_1$ describe the overscreened formation of composite pairs

$$\langle \vec{S} \cdot \psi_1 \sigma_2 \vec{\sigma} \psi_1 \rangle \propto V_1 \Delta_2 - V_2 \Delta_1$$

Decoupling the interactions

- Two channels:

$$H_I \rightarrow \left[[V_1 f^\dagger + (\Delta_1 f \sigma_2)] \psi_1 + \text{H.c} \right] + N \left(\frac{V_1 V_1 + \Delta_1 \Delta_1}{J_1} \right) \\ + \left[[V_2 f^\dagger + (\Delta_2 f \sigma_2)] \psi_2 + \text{H.c} \right] + N \left(\frac{\bar{V}_2 V_2 + \bar{\Delta}_2 \Delta_2}{J_2} \right)$$


Gauge fixing does not remove the pairing

$$\rightarrow \left[[V_1 f^\dagger \psi_1 + (\Delta_2 f \sigma_2) \psi_2] + \text{H.c} \right] + N \left(\frac{V_1 V_1}{J_1} + \frac{\Delta_2 \Delta_2}{J_2} \right)$$

The gauge-invariant cross term $V_1 \Delta_2 - V_2 \Delta_1$ describe the overscreened formation of composite pairs

$$\langle \vec{S} \cdot \psi_1 \sigma_2 \vec{\sigma} \psi_1 \rangle \propto V_1 \Delta_2 - V_2 \Delta_1$$

PC, Kee, Andrei & Tsvelik. (97)



$$\Delta_{eff}(k) \sim V_1 \Delta_2 \gamma_1(k) \gamma_2(k)$$

“resonant Andreev scattering.”

Crystal Fields determine the gap symmetry

Crystal Fields determine the gap symmetry

The gauge-invariant cross term $V_1\Delta_2 - V_2\Delta_1$ describe the overscreened formation of composite pairs


$$\langle \vec{S} \cdot \psi_1 \sigma_2 \vec{\sigma} \psi_1 \rangle \propto V_1\Delta_2 - V_2\Delta_1$$

PC, Kee, Andrei & Tsvelik. (97)

Crystal Fields determine the gap symmetry

The gauge-invariant cross term $V_1\Delta_2 - V_2\Delta_1$ describe the overscreened formation of composite pairs

$$\langle \vec{S} \cdot \psi_1 \sigma_2 \vec{\sigma} \psi_1 \rangle \propto V_1\Delta_2 - V_2\Delta_1$$




$$\Delta_{eff}(k) \sim V_1\Delta_2\gamma_1(k)\gamma_2(k)$$


Crystal Fields determine the gap symmetry

The gauge-invariant cross term $V_1\Delta_2 - V_2\Delta_1$ describe the overscreened formation of composite pairs

$$\langle \vec{S} \cdot \psi_1 \sigma_2 \vec{\sigma} \psi_1 \rangle \propto V_1\Delta_2 - V_2\Delta_1$$




(a)





$$\Gamma_7^{\pm} \otimes \Gamma_6$$



(b)



$$\Gamma_7^+ \otimes \Gamma_7^-$$



$$\Delta_{eff}(k) \sim V_1\Delta_2\gamma_1(k)\gamma_2(k)$$





Large N Kondo lattice:
overscreening drives pairing.

Confirmed:
Cox, Pang, Jarell (96)
PC, Kee, Andrei, Tsvelik (98)






Large N Kondo lattice:

overscreening drives pairing.




Confirmed:

Cox, Pang, Jarell (96)


PC, Kee, Andrei, Tsvelik (98)



Curro et al, Nature, 434,622 (2005).






Curro et al, Nature, 434,622 (2005).






Conclusions

Conclusions






Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.






Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.
- $1/N$ expansion for heavy fermion superconductivity






Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.
- $1/N$ expansion for heavy fermion superconductivity
- Predicts overscreening will lead to composite pairing and a new type of Andreev scattering.






Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.
- $1/N$ expansion for heavy fermion superconductivity
- Predicts overscreening will lead to composite pairing and a new type of Andreev scattering.
- enhancement of $1/T_1$ above T_c .



Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.
- $1/N$ expansion for heavy fermion superconductivity
- Predicts overscreening will lead to composite pairing and a new type of Andreev scattering.
- enhancement of $1/T_1$ above T_c .
- Enhancement of Andreev scattering above predictions of BTK



Conclusions

- Motivated by new superconductors, new method for preserving symplectic symmetry of spin in $1/N$ expansions.
- $1/N$ expansion for heavy fermion superconductivity
- Predicts overscreening will lead to composite pairing and a new type of Andreev scattering.
- enhancement of $1/T_1$ above T_c .
- Enhancement of Andreev scattering above predictions of BTK
- NpAl_2Pd_5 and PuCoGa_5 have identical crystal symmetry. Possibility that Pu doping will enhance T_c in NpPd_2Al_5 .

